

## RESEARCH ARTICLE

# Work-related Musculoskeletal Disorder Symptoms among Computer User Workers of Ethiopian Roads Authority in Addis Ababa, Ethiopia

Belay Negassa Gondol<sup>1\*</sup> and Adane Tesfaye Anbese<sup>1</sup>

Received: 30 December 2022

Accepted: 29 May 2023

DOI:10.20372/ajhsm.v02i01.01

Published: 21 June 2023



**Suggested Citation:** Negassa B.G. and Tesfaye A.A., Work-related Musculoskeletal Disorder Symptoms among Computer User Workers of Ethiopian Roads Authority in Addis Ababa, Ethiopia. *Afri. J. Heal. Sci. Med.*; 2023, 02(01).

**Copyright:** ©2023 Dilla University. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

## Abstract

**Background:** Computer related health problems and ergonomic disorders are resulted from prolonged computer usage in a maladaptive manner. At least 10 million new cases of computer related human health risks occur each year; posing in reduced life quality and work productivity. This study aims to describe the prevalence of work-related musculoskeletal disorder symptoms among computer user workers of Ethiopian Roads Authority.

**Methods:** Two hundred thirty three workers were requested for the study from all ten districts of Ethiopian Roads Authority from January 2018 to February 2018. A cross-sectional study was conducted using self-administered questionnaires to collect socio-demographic data, symptoms of musculoskeletal disorders, information regarding ergonomic and computer work station conditions. Chi-square test was used to determine the association between the variables. The odds ratio was calculated using binary logistic regressions. The researcher declared the level of statistical significance at p-value <0.05

**Result:** A total of 233 workers were included in this study from Ethiopian Roads Authority; 61% of the respondents were males, whereas; 39% were females. The 12-months prevalence of work related musculoskeletal disorder symptoms among computer users in this study population was 71.2%. The most frequently self-reported symptoms were back pain (46.4%), headache (39.9%), neck pain (31.3%), and wrist pain (12%) among Ethiopian Roads Authority workers ( $p < 0.05$ ). Computing hours per day was significantly associated with the presence of musculoskeletal disorder symptoms.

**Conclusion:** Ethiopian Roads Authority computer user workers had a high prevalence of work-related musculoskeletal disorders. Daily computer using hours was the most single statistically significant risk factor in this study. There is a need to increase the corrective measures that to be implemented to reduce the impact of computer related symptoms of musculoskeletal disorders.

**Keywords:** Musculoskeletal symptoms, Computer user workers, Ergonomic training, Ethiopian Roads Authority, Work-related musculoskeletal disorders

\*Correspondence: [belaynegasa13@gmail.com](mailto:belaynegasa13@gmail.com)

<sup>1</sup>School of Public Health, College of Medicine and Health Sciences, Dilla University, Dilla, Ethiopia.

Full list of author information is available at the end of the article

## Background

In today's age, computer has become a common item; its usage become an integral part of daily life [1]. Computer is changing the work environment by simplifying a number of related tasks across many organization. It has increased the work efficiency and communications and has opened access to information like never before [2]. Huge numbers of people use computers excessively and intensively starting from official work to playing video games [3].

Continuous use of computers for long hours [4] in maladaptive manner found to have severe problems; even for few hours per day usage [5], causes various health illness. Scholars [6] have identified health risks developed from usage of computer for three hours per day such as Occupational Overuse Syndrome (OOS), Computer Vision Syndrome (CVS), Musculoskeletal Symptoms (MSSs): low back pain, neck pain, wrist pain; headaches and psychosocial stress. These risks are directly related to prolonged sitting in front of computer screens with poor ergonomic practices [7,8].

Mashige *et al.* [9] and Ranganatha S C *et al.* [10] have listed posture-related symptoms as neck pain, shoulder pain, back pain, wrist pain, knee pain which are collectively named as musculoskeletal symptoms (MSSs).

Work-related musculoskeletal symptoms can occur as a result of inappropriate working conditions and poor working habits. It might be associated with maladapted monotonous use of computers that attributed to poor ergonomic practices [11]. Musculoskeletal symptoms can cause musculoskeletal disorder considered as one of the contemporary health issues. It comprises two percent of global disease burden and the second highest volume of years lived with disability [1,12]; existed in 22% of patients [13]. Part of MSDs categorized under musculoskeletal symptoms includes: headache, neck and back pain, and shoulder, wrist, and finger discomfort

[13,14]. These symptoms are well associated with improper placement of computer screen which lead to muscles sprain; affecting head and neck postures when working at a computer [15,16].

Most evidences show that, work-related musculoskeletal disorders have been called the foremost occupational hazard of the 21<sup>st</sup> century [17]. The studies have identified that 64% to 90% of computer users have health problems [18]. Study of 2014 in US shown that, on average, (45 to 70) million people spend an hour staring into a computer screen. But, recently about 143 million of workers use a computer daily [19] in U.S.A alone and 90% of whom who use three to four hours per day developed computer related illnesses [20]. Global records also estimated that nearly 60 million people suffer [6] and other study estimates that at least 10 million new cases of computer-related human health risks were reported each year [21].

Work-related musculoskeletal disorders (WMSDs) lead to economic costs and affect organizational effectiveness and efficiency. Excessive use of technology [22]; duration of occupation [23]; lack of ergonomic training for workers to implement safe practices at their work place [24] have led to increase prevalence of WMSDs. Since MSSs places an unusual strain on workers physical well-being, it decreases the quality of life [25], reduces employees' effectiveness, and significantly causes a resultant loss of productivity [26].

So, MSDs can be considered as a significant rising non-communicable disease having the capacity of global public health threat unless serious attention is paid to it [27].

Many studies have been conducted to address questions concerning safety and health for computer users. Most of these studies reported prevalence of WMSDs ranged from 40-73.3% [13,28-31]. However, they were centered in Western and Middle East regions focusing on assess-

ing knowledge, practice and magnitude of eye related problems of computer usage targeted on university communities as their study population [32-41]. But, every individual using computer for longer time without rest at their office or home can develop work-related MSSs or WMSDs [28-31].

As far as the knowledge of the authors concerned, there is limited studies related to musculoskeletal disorder documented in Africa including Ethiopia. Few studies were conducted in limited states or districts: Debre Tabor and Gonder in Ethiopia [42,43]; Abuja in Nigeria [44] assessed the prevalence of CVS among computer users. Academic and financial institutions were the subject of a few studies conducted in Ethiopia [42,43,45] to assess computer-related health issues. These studies do not go far enough to investigate the existence of computer-related health risks and predictor variables across various groups of computer users.

As a result, computer users in other broad representative federal organizations, such as the Ethiopian Roads Authority, which employs a large number of people with varying levels of computer usage, should be included in such studies. Furthermore, no single study has attempted to determine the magnitude of MSDs and its related factors among Ethiopian computer users. Posture related symptoms which are also a growing, but neglected health risks among computer users was excluded. Also evidences show that work-related MSSs lead to complicated MSDs. Therefore, the aim of this study was to determine the prevalence of work related symptoms of MSDs and their associated factors among Ethiopian Roads Authority computer users.

## Methods and Materials

### Study Design and Period

A cross-sectional study design was used from January 2018 to February 2018.

### Study Area

The study was conducted in Addis Ababa, the capital city of Ethiopia. The government institution at which this study conducted was Ethiopian Roads Authority, which consist about 34 departments (work units) with a total of 2928 workers as ERA's census of April, 2017 [47].

### Source and Study Population

All computer users who worked in Ethiopian Roads Authority were the source population, whereas all workers who were using computer in their day-to-day working activities for at least twelve months were taken as study population. Any type of duties carried out by employees was considered.

### Eligibility Criteria

The research included all ERA computer users, with the exception of those who had used a computer for fewer than twelve months. ERA employees who had a history of known musculoskeletal disorders prior to starting computer use were also ineligible.

### Sample Size Determination

The sample size (n) was first estimated using the single population proportion formula with the following assumptions: a margin of error (D) of 5%, a proportion (P) of health risk in frequent computer users 73.3 percent [21], and a confidence interval of 95 percent;  $Z_{\alpha/2}=1.96$

$$n = \frac{(Z_{\alpha/2})^2 p(1-p)}{D^2}$$

$$n = \frac{(1.96)^2 (0.73)(1-0.73)}{(0.05)^2} = 301$$

$$n_f = \frac{n}{1 + \frac{n}{N}} = \frac{301}{1 + \frac{301}{2928}} = 272$$

### Sampling Procedure

A pre-determined sample size was allocated to 34 work units of ERA offices. A random sampling technique was used to select participants. Then, from each selected office, study subjects were

selected proportionally to their size by random sampling technique.

### Operational Definition

*Musculoskeletal Disorders, MSDs:* will be considered if the respondents have experienced a variety of symptoms such as discomfort in the neck, pain in the shoulders, elbows, hands, fingers, hips and knees as a result of repetitive movements, doing work in awkward postures and static postures while prolonged seating at works [28].

*Computer user workers:* employees those using computer for their day-to-day life activities to perform different tasks [42] at least for twelve months and who were staff at ERA.

*Ergonomically nonadjustable chairs:* Non-ergonomic chairs usually have a fixed and non-adjustable backrest, which means you can't adjust it to suit your needs [23].

### Data Collection Procedure

Self-administered data collection tools were used to capture information related to the problem under investigation from the study subjects. The tool was initially designed in English language and translated into Amharic and back translated into English language for data entry and analysis. The purpose and objectives of the study was debriefed to the study subject before consent. The questionnaire was distributed to each working stations and completeness of the tool was cross-checked before collecting back from the respondents.

### Data Analysis

Data was entered to Excel spread sheet and imported to predictive analytics software 20 version software for statistical analysis (IBM SPSS Statistics 20). The prevalence of computer re-

lated disorders was expressed as percentage. Chi-square test was used to determine the association between the variables. The odds ratio (OR) was determined using logistic regressions. The significance level was considered as  $P < 0.05$ .

### Ethical Approval

The study was approved by the Ethics committee of the Ethiopian Roads Authority and therefore been performed in accordance with the ethical standards.

## Results

### Socio-demographic Characteristics

A total of 233 computer user ERA workers who met the criteria were included in the study with a response rate of 85.7%. This is due to incorrect response of 39 participants. Male participants were 142 (61%) and 91 (39%) of them were females. Majority of the study subjects (40.4% were between the age of 21-30 years followed by 33.3% of the study subjects fall between 31-40 years of age. One hundred forty one study subjects were having masters level of educational background and the remaining study participants (25.3%) had college diploma. The majority of the respondents (42.5% were engineer by profession.

Six out of ten study subjects (59.7%) have been reported previous work experience ranging from 1 to 5 years but 24% of the respondents had 6 to 10 years of previous work experiences. Only 16.3% of the study participants have previous work experience above 10 years. Nearly 75% of the study subjects were using computer for more than 6 hours per day. Hence, this is in agreement with the normal office hours in Ethiopia. About 14.2% were using computer for 3 to 5 hours a day and the rest 11.2% were using for less than 2 hours a day.

**Table 1** Socio-demographic Characteristics of Computer User Workers of ERA in Addis Ababa, 2018

Variables	All(N=233)	Does MSD Symptoms Exist?	
		Yes	No
Gender			
Male	142(61%)	101(71.1%)	41(28.9%)
Female	91(39%)	65(71.4%)	26(28.6%)
Age			
21-30 years	96(41.2%)	66(39.8%)	30(44.8%)
31-40 years	77(33.3%)	60(36.1%)	17(25.4%)
41-50 years	40(17.2%)	25(15.1%)	15(22.4%)
51-60 years	20(8.6%)	15(9.0%)	5(7.5%)
Education level			
Diploma	59(25.3%)	37(22.3%)	22(32.8%)
BA/BSc degree	141(60.5%)	105(63.3%)	36(53.7%)
MA/MSc degree	33(14.2%)	24(14.5%)	9(13.4%)
Occupation type			
Engineer	99(42.5%)	78(47.0%)	21(31.3%)
Finance officer	38(16.3%)	23(13.9%)	15(22.4%)
Human resource officer	35(15.0%)	26(15.7%)	9(13.4%)
Office manager/secretary	38(16.3%)	29(17.5%)	9(13.4%)
Others	23(9.9%)	10(6.0%)	13(19.4%)
Service year on current job			
1-5 years	139(59.7%)	103(62.0%)	36(53.7%)
6-10 years	56(24.0%)	36(21.7%)	20(29.9%)
11-15 years	16(6.9%)	13(7.8%)	3(4.5%)
Above 15 years	22(9.4%)	14(8.4%)	8(11.9%)
Ergonomic training			
Yes	29(12.4%)	17(10.2%)	12(17.9%)
No	204(87.6%)	149(89.8%)	55(82.1%)
Daily computer usage time			
More than 6 hours	174 (74.7%)	141 (84.9%)	33(49.3%)
3 to 5 hours	33(14.2%)	19(11.4%)	14(20.9%)
1 to 2 hours	26(11.2%)	6(3.6%)	20(29.9%)
Adjusting screen			
Yes	203(87.1%)	151(91.0%)	52(77.6%)
No	30(12.9%)	15(9.0%)	15(22.4%)

### Prevalence of MSD Symptoms

One hundred sixteen employees were reported a history of one or more symptoms of musculoskeletal disorder. But a 12 months prevalence of MSD among the study subjects was found to be 71.2% which is 166 out of 233. The most commonly reported work-related musculoskeletal disorder (WMSD) complaint was back pain

(46.4%); followed by headache (39.9%), neck pain (31.3%) and Wrist pain (12%) among musculoskeletal symptoms, MSSs (Figure 1). The prevalence of symptoms ranged from the least frequently reported (4.3%) for shoulder pain to (46.4%) for back pain among ERA workers. Almost there was no difference in cumulative prevalence of WMSD symptoms among male (71.1%) and female (71.4%) workers ( $p > 0.05$ ).




Figure 1 Musculoskeletal Symptoms of Computer User Workers in ERA

Table 2 One Year Prevalence of MSD Symptoms among Computer User Workers of Ethiopian Roads Authority, 2018

Symptoms	All prevalence %	Male prevalence %	Female prevalence %	<i>p-value</i>
Does MSSs or MSDs exist?				
Yes	166(71.2%)	101(71.1%)	65(71.4%)	.54
No	67(28.8%)	41(28.9%)	26(28.6%)	
Musculoskeletal symptoms reported				
Headache	93(39.9%)	60(42.3%)	33(36.3%)	.41
Neck pain	73(31.3%)	49(34.5%)	24(26.4%)	.20
Elbow pain	20(8.6%)	12(8.5%)	8(8.8%)	.55
Wrist pain	28(12.0%)	10(7.0%)	18(19.8%)	.004*
Shoulder pain	10(4.3%)	8(5.6%)	2(2.2%)	.33
Back pain	108(46.4%)	70(49.3%)	38(41.8%)	.29
Thigh pain	15(6.4%)	13(9.2%)	2(2.2%)	.051
Knee pain	18(7.7%)	12(8.5%)	6(6.6%)	.63

**Workstation environs and Ergonomics related Problems Triggers MSSs**

The ergonomic hazards were part of the assessment conducted during the survey. Almost more than half of the respondents 120 (51.5%) were used ergonomically nonadjustable chairs. More than half (54.1%) of the participants usually used desktop computers. Nearly half of them 114 (48.9%) were reported inadequate free space

near to their computer work station.

About 12.9% of respondents reported that they were not adjusting their computer screen. On the other hand, 87.6% of the participants did not attend any ergonomic training related to computer usage (Table 1 and 2).

**Table 3** Working Station Environs and Percentage of MSSs among Computer Users in Ethiopian Roads Authority (n = 233), 2018

Symptoms of MSD	Variables (working station environment)									
	Computer type used				Using adjustable chair			Free space availability		
	Dt	Dt & Lt	Lt	P-value	Yes	No	P-value	Yes	No	P-value
Headache	34.9	47.7	38.1	.17	42.5	37.5	.43	47.9	31.6	.01*
Neck pain	30.2	31.4	38.1	.78	31.0	69.0	1.0	31.1	68.9	.52
Elbow pain	4.8	11.6	19.0	.04	8.0	92.0	.81	7.6	92.4	.36
Wrist pain	8.1	15.1	9.5	.31	11.5	88.5	.84	14.3	85.7	.18
Shoulder pain	2.4	8.1	0.0	.10	7.1	92.9	.054	5.9	94.1	.33
Back pain	46.0	48.8	38.1	.68	47.8	52.2	.69	47.9	52.1	.69
Thigh pain	4.8	9.3	4.8	.48	5.3	94.7	.59	5.9	94.1	.79
Knee pain	9.5	5.8	4.8	.61	7.1	92.9	.80	8.4	91.6	.81

Dt=Desktop; Lt=Laptop

**Factors Associated with MSDs Symptoms**

Job category, ergonomic training, computing hours per day and adjusting computer screening were candidate variable with the occurrence of MSSs ( $p < 0.25$ ). However, among all candidate variables only patterns of computer usage per day was significantly associated with occurrence of MSDs ( $p < 0.05$ ).

**Results of the Logistic Regression Analysis**

From binary logistic regression analysis: computing hours per day was significantly associated with the presence of MSD symptoms. When compared to workers who use the computer for more than 6 hours a day, using the computer for 1 to 2 hours a day substantially reduced the incidence of MSD symptoms by 16.4 times (AOR=16.401; 95% CI: 5.371-50.086).

**Table 4** Factors Associated with MSD Symptoms among Computer User ERA Workers (n=233), 2018

Variables	COR (95% CI)	AOR (95% CI)	P-value
Job category			
Engineer <sup>®</sup>			
Finance and accountant	2.42 (1.08-5.44)*	2.16 (0.90-5.21)	0.085
Human resource officer	1.29 (0.52-3.16)	1.10 (0.39-3.07)	0.861
Office manager/secretary	1.15 (0.47-2.81)	0.47 (0.16-1.38)	0.168
Others	4.83 (1.86-12.55)*	2.89 (0.34-24.94)	0.334
Ergonomic training			
Yes <sup>®</sup>			
No	0.52 (0.24-1.17)	0.84 (0.31-2.26)	0.730
Computer usage per day			
More than 6 hours <sup>®</sup>			
3 to 5 hours	3.15 (1.43-6.92)*	3.814 (1.65-8.84)	0.002
1 to 2 hours	14.24 (5.30-38.25)*	16.40 (5.37-50.09)	0.000
Adjusting computer screen			
Yes <sup>®</sup>			
No	2.90 (1.33-6.35)*	0.98 (0.15-6.23)	0.983

Note: <sup>®</sup> = Reference; \* = indicates candidate variables for multivariable

## Discussion

In this study, 71.2% of the employee had one or more than one symptoms of MSDs revealing the highest health related risk associated with computer utilization. This finding is higher than studies reported prevalence of MSD symptoms ranged from 40-70% [22,24,33-35,42-45].

Higher prevalence found in this study is probably due to higher sample size with the reference to similar previous studies. Then again, most respondents in this study were engineers, who were experienced on daily usage of computers for long period of time as evidenced by Logaraj *et al.* [34]. Hence, these inconsistencies might be a possible justification for the determined higher prevalence in present study.

The main work-related musculoskeletal disorder symptoms of this study were back pain (46.4%), headache (39.9%), neck pain (31.3%) and wrist

pain (12%) among ERA workers. This is aligned with a study done by other scholars [34,40].

There is no significant sex difference in the overall prevalence of MSDs among male (71.1%) and female (71.4%) employees (p-value >0.05). But, in particular wrist (p = 0.004) pain were observed symptoms of MSDs associated to gender (p <0.05) and more prevalent among female participants which is similar to Pandey *et al.* [12].

The possible explanation for the similarity might be due to the proportion of male and female employees. In this study, 41.8% of the employee were female and office secretary to have a higher chance of developing wrist/ fingers pain as a result of frequent computer use through out their office hours. Strengthening this fact, Dessie *et al* found that secretaries were significantly impacted by MSDs compared to other workers [42].

Out of all, 233, respondents met the criteria

and participated in this study, the majority 96 (40.4%) of them belonged to age category of 21-30 years old followed by age groups between 31-40 years 77 (33.3%). No significant association was found between the ages of workers with MSDs development in present study. Nevertheless, it might be revealed that most computer based work was dominated by younger generation. This is in line with other study conducted by Abudawood *et al.* in King Abdulaziz University, Jeddah, Saudi Arabia [40]. The current finding disagree with the study by Alemayehu *et al.* [45] who reported that older aged were at higher risk of developing computer related health problems within their study population.

Daily computer using time was the most statistically significant risk factor in this study. Workers who used computers for 3 to 5 hrs per day (AOR: 3.8; 95% CI=1.6-8.8, p=0.002) were 3.8 times less likely to develop MSD symptoms as compared to those who used computers more than 6 hours per day. On the other hand, computer user workers using for 1 to 2 hours per day (AOR: 16.4; 95% CI=5.37-50.08, p=0.000) were 16.2 times less likely to develop symptoms of MSDs comparing to those who used more than 2 hours per day. This finding is in agreement with numbers of the previous findings [22,23,36,38].

Moreover, Noreen *et al.* [22] and Logaraj *et al.* [34] reported that users who spent continuously without rest more than four hours were significantly at higher risk of developing computer related health problems than who spent less than four hours. Other similar studies also shown that the longer the time spent on computer, the more prevalent and extent are risks of MSDs symptoms appreciably [4,24]. Additionally, Logaraj *et al.* [34] observed that the longer the duration, the longer the complaint last even after work. Hence, either reducing daily exposure time spent on computer or taking mini breaks [23] are important to prevent and control MSDs.

Neck pain was more prevalent among laptop

users (40%) (table 3), this might be resulted from view distance which was revealed by scholars in previous studies [33,46]. Users who viewed computers at a distance of less than arm and forearm length (> 50 cm), resulted in more symptoms significantly [46]. Shantakumari *et al.* [33] added that the prevalence of headache decreased in students who viewed the screen at a distance more than 50 cm.

Most chairs used by computer users in properly designed computer facilities and institutions have adjustments to make them comfortable to sit on and therefore preventing back pains [23]. However, almost 5 out of 10 (51.5%) of ERA workers were sitting on non-adjustable chair in present study. This revealed that only 48.5% of the workers used adjustable chair which was inconsistent with Logaraj *et al.* [34] who reported that 61.5% of respondents in their study used chairs with adjustable backrest while working on their computer.

Despite it was non-significant in this study, sitting on inappropriate chair in front of computer screen cause muscle stiffness, headache, and back pain as muscles and tendons become inflamed due to prolonged sitting [12]. Strengthening this, evidence explaining that musculoskeletal symptoms are well related to improper seating posture and placement of the screen [16]. Additionally, almost half of the study subjects (48.9%) were complaining the importance of the availability of free space near to their work station.

Although, this is not significant in present study, it may be one factor that triggers for 46.4% back; 39.9% headache; 31.3% neck; 12% wrist, 8.6% elbow; 7.7% knee, 6.4% thigh and 43% shoulder pains among study participants. That's why, American Optometric Association recommend that proper ergonomic design and adjustment of computer in an adequate workstation can increase productivity and workers comfort by decreasing the demands of the task [15].

## Limitation of study

Self-reported symptoms of MSDs were considered by excluding ergonomic examinations using cross-sectional study is the limitation of the study. Other ergonomic parameters like workstation furniture and distance of computer screen from eye contact might be related to MSSs were not taken into the account.

## Conclusion

Ethiopian Roads Authority computer user workers had a high prevalence of work-related musculoskeletal disorder symptoms. Computing hours per day was significantly associated with the presence of musculoskeletal disorder symptoms. Multi-programmed approaches; decreasing the number of hours per day at the computer screen, increasing frequent rest (mini breaks) with motivation for exercises (to stand, stretch, and move around) at workplace and proper positioning is needed to prevent Work-related Musculoskeletal Disorder Symptoms (WMSDs). Ergonomic interventions should be considered to prevent the computer related health problems.

## Assertions

### Abbreviations

CVS	Computer Vision Syndrome
CI	Confidence Interval
ERA	Ethiopian Roads Authority
MSSs	Muscular Skeletal Symptoms
MSDs	Musculoskeletal Disorders
WMSDs	Work-related Musculoskeletal Disorders
OOS	Occupational Overuse Syndrome
SPSS	Statistical Package for Social Sciences

### Acknowledgments

The author would like to express his gratitude to ERA's principals and employees for their willingness to participate in this research. Thank you to Mrs. Birhan Hagos, Mr. Dawit Solomon, and Amha Hailu for their assistance in revising the questionnaire.

**Consent for publication** - Not applicable

### Availability of data and materials

The datasets underlying the study are available from the corresponding author up on request.

## Competing interests

The author declares that there is no competing interest.

## Author's detail

<sup>1</sup>School of Public Health, College of Medicine and Health Sciences, Dilla University, Dilla, Ethiopia.

## References

1. Mani, K., *Ergonomics Education for Office Computer Workers: An Evidence-Based Strategy, in Anatomy, Posture, Prevalence, Pain, Treatment and Interventions of Musculoskeletal Disorders*. 2018, IntechOpen.
2. Gangamma, M.P., Poonam, and M. Rajagopala, *A clinical study on "Computer vision syndrome" and its management with Triphala eye drops and Saptamrita Lauha*. Ayu, 2010. **31**(2): p. 236-9.
3. Randolph, S.A., *Computer Vision Syndrome*. Workplace Health & Safety, 2017.
4. Sultan H. Al Rashidi and H. Alhumaidan, *Computer vision syndrome prevalence, knowledge and associated factors among Saudi Arabia University Students: Is it a serious problem*. International Journal of Health Sciences, 2017. **11**(5).
5. Kumar, B.S., *A Study to Evaluate the Knowledge Regarding Computer Vision Syndrome among Medical Students*. Biomedical & Pharmacology Journal, 2020. **13**(1): p. 469-473.
6. Sen A. and S. Richardson, *A Study of Computer-Related Upper Limb Discomfort and Computer Vision Syndrome*. J. Human Ergol, 2007. **36**: p. 45-50.
7. Mohsen Soroush and H. Hassani, *Musculoskeletal complaints associated with computer use and its ergonomic risks for office workers of a medical sciences university in Tehran*. Annals of Military & Health Sciences Research, 2015. **13**(1): p. 2-6.
8. *Guidelines for using computers – preventing and managing discomfort, pain and injury*, D.o. Labor, Editor. 2010, Accident Compensation Corporation (ACC): New Zealand.
9. Khathutshelo Percy Mashige, N. Rampersad, and O. Oduntan, *A study of ergonomic factors leading to computer vision syndrome among computer users*. Ergonomics SA, 2013. **25**(1).
10. Ranganatha S C and S. Jailkhani, *Prevalence and Associated Risk Factors of Computer Vision Syndrome among the Computer Science Students of an Engineering College of Bengaluru- A Cross-Sectional Study*. Galore International Journal of

- Health Sciences and Research, 2019. **4**(3): p. 10-15.
11. Soumya Harapanahalli Venkatesh, Anita T Girish, Shashikala, Praveen Kulkarni, Snigdha Mannava, Rajendra Rajarathnam, *A Study of Computer Vision Syndrome at the Workplace -Prevalence and Causative Factors*. International Journal of Contemporary Medical Research, 2016. **3**(8): p. 2375-2377.
  12. Rahul Pandey, Shikhar Gaur, Rakesh Kumar, Narendra Kotwal, Sanjeev Kumar, *Curse of the technology-computer related musculoskeletal disorders and vision syndrome: a study*. Int J Res Med Sci., 2020. **8**(2): p. 661-666.
  13. Tasneem Borhany, Erum Shahid, Wasim Ahmed Siddique, Hussain Ali, *Musculoskeletal problems in frequent computer and internet users*. Journal of Family Medicine and Primary Care, 2018. **7**(2): p. 337-339.
  14. Lurati, A.R., *Computer Vision Syndrome: Implications for the Occupational Health Nurse*. Workplace Health & Safety, 2018. **66**(2): p. 56-60.
  15. American Optometric Association, *The Effects of Computer Use on Eye Health and Vision*. 1997, American Optometric Association: St. Louis, MO, USA.
  16. Loh KY and R. SC, *Understanding and Preventing Computer Vision Syndrome*. Malaysian Family Physician, 2008. **3**(3): p. 128-130.
  17. Shweta Chauhan, Renu Dhasmana, and A. Raj, *Knowledge, Awareness and Practice of CVS in Digital Device Users*. Sudanese Journal of Ophthalmology, 2018. **10**(1).
  18. Rosenfiel, M., *Computer vision syndrome: a review of ocular causes and potential treatments*. Ophthalmic Physiol Opt, 2011. **31**: p. 502-515.
  19. Saad Abbas Aldawsari, Abdullah Ayidh Alotaibi, Khalid Mohamed Alabdulwahhab, Elsadig Yousif Mohamed, Sharafeldin Abdelmajid, *Knowledge, attitudes and practices of faculty members' about computer vision syndrome, Majmaah University, Saudi Arabia*. Int J Community Med Public Health, 2018. **5**(9): p. 3801-3806.
  20. Hafiz Muhammad Junaid Hassan, Sarah Ehsan, and H.S. Arshad, *Frequency of Computer Vision Syndrome & Ergonomic Practices among Computer Engineering Students*. International Journal of Science and Research, 2016. **5**(5): p. 121-125.
  21. Manish A Prasad, Vasant Wagh, and A. Mudey, *Study of Prevalence of Health Problems among Computer Professionals in Selected Information Technology (IT) Company in Nagpur District of Central India*. Innovative Journal of Medical and Health Science, 2014. **4**(3): p. 96 - 98.
  22. Khola Noreen, Zunaira Batool, Tehreem Fatima, Tahira Zamir, *Prevalence of Computer Vision Syndrome and Its Associated Risk Factors among Under Graduate Medical Students*. Pak J Ophthalmol, 2016. **32**(3): p. 140-146.
  23. Mohamed Sherif Sirajudeen and S.S.M. Siddik, *Knowledge of Computer Ergonomics among Computer Science Engineering and Information Technology Students in Karnataka, India*. Asian Journal of Pharmaceutical Research and Health Care, 2017. **9**(2): p. 64-70.
  24. Ranasinghe P, W. S. Wathurapatha, Y. S. Perera, D. A. Lamabadusuriya, S. Kulatunga, N. Jayawardana, P. Katulanda, *Computer vision syndrome among computer office workers in a developing country: an evaluation of prevalence and risk factors*. BMC Research Notes, 2016. **9**(150).
  25. Mrinal Ranjan Srivastava, Anu Chandra, and S. Choudhary, *Computer Vision Syndrome: New Age Eye Malady-A Short Review*. Acta Scientific Medical Sciences, 2020. **4**(4): p. 10-18.
  26. Akinbinu T. R and M.Y. J, *Impact of computer technology on health: Computer Vision Syndrome (CVS)*. academic Journals, 2014. **5**(3): p. 20-30.
  27. Agbonlahor, *Prevalence and knowledge of Computer Vision Syndrome (CVS) among the Working Class Adults in F.C.T. Nigeria*. Journal of the Nigerian Optometric Association, 2019. **21**(1): p. 49-60.
  28. Mohsen Soroush MD and H.H. MS, *Musculoskeletal complaints associated with computer use and its ergonomic risks for office workers of a medical sciences university in Tehran*. Annals of Military & Health Sciences Research, 2015. **13**(1).
  29. Ehsanollah Habibi, Zahra Mohammadi, and A.G. Sartang, *Ergonomic assessment of musculoskeletal disorders risk among the computer users by Rapid Upper Limb Assessment method*. Int J Env Health Eng, 2016. **5**(2): p. 1-4.
  30. Nabila Chowdhury, Fereydoun Aghazadeh, and M. Amini, *Ergonomic assessment of working postures for the design of university computer workstations*. Occupational Ergonomics, 2017. **13**: p. 37-46.
  31. Mazhar Ali Bhutto, Arsalan Abdullah, Syed Asadullah Arslan, M Sarfraz Khan, Muhammad Khan Bugti, Zeb Jehan Rana, *Prevalence of Neck Pain in Relation to Gender, Posture and Ergonomics in Computer Users*. Acta Scientific Orthopaedics, 2019. **2**(10): p. 02-06.
  32. Reddy SC, Low CK, Lim YP, Low LL, Mardina F, Nursaleha MP, *Computer vision syndrome: a*

- study of knowledge and practices in university students.* Nepal J Ophthalmol, 2013. **5**(10): p. 161-168.
33. Shantakumari N, Eldeeb R, Sreedharan J, Gopal K, *Computer Use and Vision-Related Problems Among University Students In Ajman, United Arab Emirate.* Annals of Medical and Health Sciences Research, 2014. **4**(2): p. 258-263.
  34. Logaraj M, Madhupriya V, and H. SK, *Computer Vision Syndrome and Associated Factors Among Medical and Engineering Students in Chennai.* Annals of Medical and Health Sciences Research, 2014. **4** (2): p. 179-185.
  35. Huda Zainuddin and M.M. Isa, *Effect of Human and Technology Interaction: Computer Vision Syndrome among Administrative Staff in a Public University.* International Journal of Business, Humanities and Technology, 2014. **4**(3): p. 39-44.
  36. Ashlesh Upadhyaya and R. Joshi, *An Evaluation of Knowledge and Attitude amongst Adolescents of Ahmedabad towards Computer Vision Syndrome and Effectiveness of Structured Teaching Programme.* International Journal of Health Sciences & Research, 2014. **4**(10): p. 191-198.
  37. Ranju Kharel (Sitaula) and A. Khatri, *Knowledge, Attitude and practice of Computer Vision Syndrome among medical students and its impact on ocular morbidity.* J Nepal Health Res Counc, 2018. **16**(40): p. 291-296.
  38. Mowatt, L., Gordon, C. Santosh, A. B. R. Jones, T., *Computer vision syndrome and ergonomic practices among undergraduate university students.* Int J Clin Pract, 2018. **72**(1).
  39. Precious Emmanuel Chisale, Thokozani Mzumara, and J.C. Afonne, *Knowledge, Attitude, Perception and Knowledge and Practice of Prevention Practices of Computer Vision Syndrome among Mzuzu University Academic Staff.* Journal of Eye and Vision, 2019. **2**(1:2).
  40. Ghufran A. Abudawood, Heba M. Ashi, and N.K. Almarzouki, *Computer Vision Syndrome among Undergraduate Medical Students in King Abdulaziz University, Jeddah, Saudi Arabia.* Journal of Ophthalmology, 2020: p.7.
  41. Altalhi, A., Khayyat, W. Khojah, O. Alsalmi, M. Almarzouki, H., *Computer Vision Syndrome Among Health Sciences Students in Saudi Arabia: Prevalence and Risk Factors.* Cureus, 2020. **12**(2): p. e7060.
  42. Awrajaw Dessie, Fentahun Adane, Ansha Nega, Sintayehu Daba Wami and Daniel Haile Chercos, *Computer Vision Syndrome and Associated Factors among Computer Users in Debre Tabor Town, Northwest Ethiopia.* Journal of Environmental and Public Health, 2018. **2018**: 8 pp. <https://doi.org/10.1155/2018/4107590>
  43. Natnael Lakachew Assefa, Dawit Zenebe Wel-demichael, Haile Woretaw Alemu, Dereje Hayilu Anbesse, *Prevalence and associated factors of computer vision syndrome among bank workers in Gondar City, northwest Ethiopia.* Clinical Optometry, 2017. **9**: p. 67-76.
  44. Akinbinu T.R. and Mashalla Y.J., *Knowledge of computer vision syndrome among computer users in the workplace in Abuja, Nigeria.* Journal of Physiology and Pathophysiology, 2013. **4**(4): p. 59-63.
  45. Mekuriaw Alemayehu, Ansha Nega, Eniyew Tegegne, Yohannis Mule, *Prevalence of Self Reported Computer Vision Syndrome and Associated Factors among Secretaries and Data Processors Who are Working in University of Gondar, Ethiopia.* Journal of Biology, Agriculture and Healthcare, 2014. **4**(15).
  46. Abida Ellahia, M. Shahid Khalil, and F. Akram, *Computer users at risk: Health disorders associated with prolonged computer use.* E3 Journal of Business Management and Economics, 2011. **2**(4): p. 171-182.
  47. Ethiopian Roads Authority. [http://www.era.gov.et/web/guest/publications/-/document\\_library\\_display/kPb3/view/1069608](http://www.era.gov.et/web/guest/publications/-/document_library_display/kPb3/view/1069608), Employment Record April, 2017.