

RESEARCH ARTICLE

Essential newborn care practice and associated factors among Health professionals at rural Gedeo Zone, southern Ethiopia: Cross sectional study

Moges Mareg^{1*}, Yohannes Addisu², Getachew Nenko², and Zelalem Belayneh³

Received: 22 September 2022

Accepted: 27 April 2023

DOI:10.20372/ajhsm.v02i01.04

Published: 01 June 2023



Suggested Citation: Mareg M., Addisu Y., Nenko G., and Belayneh Z., Essential newborn care practice and associated factors among Health professionals at rural Gedeo Zone, southern Ethiopia: Cross sectional study. *Afri. J. Heal. Sci. Med*; 2023, 02(01).

Copyright: ©2023 Dilla University. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Background: The neonatal mortality (NM) accounts 43% of child mortality under 5 year in the globe, and 42% in Ethiopia. There is inadequate evidence concerning the new born care practice of health workers in rural Gedeo zone of southern Ethiopia. The main objective of this study was to assess the essential newborn care practice and associated factors among health workers at rural Gedeo zone.

Methods: An institution based cross-sectional study was conducted. A total of 233 health workers (using systematic random selection method) were participated. The EPI-info Version 3.5 for data entry and statistical package for social sciences (SPSS) version 20 software was used for data analysis. Significant association was declared by P-value of < 0.05 with a corresponding 95% confidence interval.

Result: a total of 233 medical professionals took part, with a 100% response rate. Of the 233 medical professionals, 24% had good essential newborn care practices at 95% CI (19.3, 28.9), and more than half, 76.0%, missed one or more crucial steps in the care of a new born. Essential newborn care practices were significantly correlated with older age [adjusted odds ratio (AOR) = 0.18; 95% CI (0.03, 1.12)], female sex [AOR = 3.08; 95% CI (1.07, 8.81)], supportive supervision [AOR = 2.17; 95% CI (1.12, 4.19)], attending in-service training [AOR = 4.85; 95% CI (2.33, 10.10)], and being aware of the newborn's complications [AOR = 1.98; 95% CI (1.01, 3.89)].

Conclusion: over three-fourths of medical personnel failed to perform at least one crucial newborn care technique. The results suggested that in order to improve the critical newborn care provided by the medical staff at rural health institutions in Gedeo Zone, supportive supervision, in-service training, and awareness of newborn complications should be strengthened.

Keywords: Ethiopia, Health Professionals, Neonates, Newborn care Practice

*Correspondence: metanmann@gmail.com; Tel. +251923527030

¹Department of Reproductive Health, School of Public health, Medical and Health Science College, Dilla University, Dilla, Ethiopia

Full list of author information is available at the end of the article

Background

Globally, newborn mortality accounts 45% of less than 5 year old child deaths and account for 40% of under 5 year old child deaths in Tanzania, 42% of less than 5 year old child deaths in Ethiopia [1]. Seventy five percent of all neonatal mortality (deaths in the first 28 days of life) occurs in twelve countries, six of them are in sub-Saharan Africa including Ethiopia [2,3].

Birth of newborn is the main challenge to the newborn to transfer positively from intra-uterine to extra-uterine life [4]. Newborn birth and the first few hours of life are serious periods for the advance growth and development of a newborn [5,6]. Amongst the lifespan of the infantile period, newborn period (age of newborn between 1 and 28 weeks) is very crucial as it defines the whole well-being of the child and his/her feature mature life of adult [7]. Utmost newborns are born healthy and at term (gestational age greater than 37 weeks), but the essential newborn care (ENC) they obtain in the first 60 minutes, days, and weeks of existence can define their future health status [8].

Essential new born care practice (ENCP) is an all-inclusive method intended to improve the health status of new born through interventions before gestation, at pregnancy, on and shortly after natal and in the post-delivery period. Essential newborn care protects the newborn against mortality and morbidity which comprises quick commencement and exclusive breastfeeding, warm air care (comprising early drying and jacket at birth, increasing skin-to-skin contact, hindered washing, preserving “warm chain”) and cleanliness practices (comprising cord-care and health professionals hand washing) [9,10].

Neonatal mortality is concentrated at less developed countries due to high maternal mortality, and poor essential newborn care [11,12]. In the sub-Saharan Africa countries, maternal death is the uppermost in the glob (56%) and viewing

the least advancement among the regions’ maternal and new born health for over two decades [13].

The infant and neonatal mortalities rate in Ethiopia was 48/1,000 live births and 29% based on a report of the 2016 Ethiopian Demographic Health Survey (EDHS) respectively [14], Whereas the report of 2019 mini Ethiopian demographic health survey (MEDHS) showed that the current neonatal mortality of Ethiopia increased from twenty nine to thirty percent per 1,000 live births [14].

The major cause of newborn death (less than 28 days) is neonatal asphyxia and birth trauma, which accounts 34%, prematurity 25%, sepsis and other infectious conditions 18%, congenital anomalies 10%, pneumonia 6%, other conditions 4%, tetanus 2%, diarrheal diseases 1%, injuries 1% [1]. In Ethiopia, the primary causes of newborn death (less than 28 days) are neonatal asphyxia and birth trauma, which accounts 34% and becomes higher in rural areas [3]. These conditions could be prevented through quality care (good essential newborn care practice) provided by health care professionals [2,3].

However, in Ethiopia still there is a huge gap in the implementation of essential newborn care practice, studies in different areas revealed that 48.9% and 27.23% of health professionals working in health centers had poor newborn care practice, and educational status, training status, the number of training, national guideline availability, presence of sufficient equipment, health facility type and knowledge level are some of the factors associated with ENCP [11,12]. These studies were conducted at urban areas of Ethiopia. Despite its crucial needs, limited evidence is available regarding ENCP of health professionals at rural Gedeo zone. As a result, the key objective of this study was to evaluate the ENCP and associated factors among health professionals working in rural health institutions.

Methods

Study Design and Period

A facility-based cross sectional study was employed starting from February 10th - April 30th 2017. It was conducted at Gedeo zone rural districts. Gedeo zone is one of the fourteen Zones in the Southern Nations, Nationalities, and Peoples' Region (SNNPR) of Ethiopia. The two Dilla and Yirgachefe towns are the administrative centers and the former is the Zonal administrative town.

Dilla town is situated at three hundred sixty kilo meters South East from Addis Ababa (the capital city of Ethiopia). According to the Zonal health department, there are six (Bule, Gedeb, Wonago, Kochere, Yirgachefe, and Dilla Zuria) Woredas. Gedeo Zone has one referral hospital, three primary hospitals and thirty two health centers in which ENCP is implemented. Within the four selected districts of the Zone, there are 477 health workers (63 health officers, 13 BSc nurses, 4 Bsc midwives, 68 diploma midwives, and 329 clinical nurses).

Sample size and sampling procedure

The researchers used single population proportion formula by considering the assumptions made for sample size calculation i.e. a 95% confidence interval (CI), 5% margin of error and essential newborn care practice (52.4%) based on a study done in Tigray(13). And the results become 382. Since the source of population is below 10,000 (i.e. $N=477$), the researchers used correction formula, the sample were 212. The sample size for the second objective (factors associated with ENCP) was also calculated; and the maximum sample size obtained from the correction formula was taken. By taking a ten percent of non-response rate, the final sample size became 233.

First, from a total of six districts, four districts (Bulie, Wonago, Yirgachefe and Dilla Zuriya) were randomly selected. Within the four ran-

domly selected districts, there were about 23 health facilities (consisting of 477 health professionals) in which essential newborn care practice was implemented. Then, the final sample size (i.e. 233) was allocated based on the number of health professionals to the 23 health facilities proportionally. After proportional allocation of samples, random sampling technique (i.e. systematic random sampling) was used in order to identify study participants from each health center using skipping interval (K). The skipping interval (K) was obtained by dividing the total numbers of study participants available at the health center to the number of proportionally allocated sample size for that health center. Simple random sampling method was used to select the first study participant between 1 and K . Then, the skip interval (K) value was added subsequently till the planned sample size was achieved.

Data collection instruments

An interviewer administered questionnaire was used for data collection together with direct observations checklist. The interviewer administered semi structured questionnaire was used to assess the socio-demographic characteristics, supervision status and knowledge related to ENCP. The level of ENCP was assessed using a standardized interview guide/tool having two main parts; direct observations check list and interviewer administered questionnaire [14].

The new born care practice was assessed by using observation checklist summarized based on ten important procedures. The level of practice was considered as "**good**" for participants who practiced the whole ten procedures (Immediately dry the whole body-while assessing the baby's breathing- evaluate breathing (if not birthing-call for help and start resuscitation), Cord care (clamping, tying and cutting, sterile blade after 1-3minute of birth), place neonate on mother's abdomen or kangaroo mother care (KMC), Helps to initiate immediate breastfeeding within 1

hour, introduce eye drop/tetracycline eye cream, provide Vitamin K injection, Place the newborn's ID bands on the wrist and ankle, Weigh the newborn, Vaccinate at birth(OPV0 and BCG) given correctly or "**poor**" if the respondent misses one or more of the 10 step procedures [14].

Knowledge of health professionals regarding newborn care was assessed by using 28 item questions. Respondents who answered above 75% of the questions correctly were considered as "have good knowledge", 50-75% "have fair knowledge" and less than 50% "not knowledgeable" [14].

Data Collection Procedure

First the tools were developed in English and then converted to Gede'uffa and Amharic (the commonly spoken language in the study area) and again to English to check its uniformity. Eight clinical nurses supervised by two BSc level health officers were involved in the data collection after 2 days training. Then, the tool was pre tested at Chuko health center among 5% (12 health professionals) before the actual data collection. The supervisors and data collectors were assigned at different health centers on daily basis. The data were collected by using observation checklist and interviewer administered questionnaire. The data were collected from health professionals by direct observation to see whether they perform the newborn care in line with the 10 essential newborn care procedures or not. The health professionals were aware of whether they were being observed or not during the data collection. After that the researchers interviewed them to get socio demographic, knowledge, training and supportive supervision related information.

Data quality control

Before the actual data collection, an English version questionnaire was translated to the local languages (Gedeufa and Amharic) to make it more understandable. Training was delivered

for supervisors and data collectors regarding the contents of the questionnaire, data collection procedures and what they can do if a professional is not knowledgeable for 2 days. Moreover, pretest of the questionnaire was done at Chuko health center among 5% (12 health professionals) before collecting the actual data collection. After the pretest, the expressions of some questions were modified using simple words.

Data processing and management

First, the data were intensively checked for its comprehensiveness and uniformity. After checking, the data were entered to EPI-info-version 3.5 and exported to statistical package for social science (SPSS) statistical software version 20 for analysis. Descriptive statistics was used to quantify the level of ENCP and the characteristics of the study participants. Both Bi-variable and multivariable logistic regression analysis were employed to recognize the factors related with ENCP. After Bi-variable analysis, the variables with a P-value < 0.25 were considered as candidate for multivariable logistic regression analysis. In multivariable logistic regression analysis, P-value of < 0.05 was considered as statistically significant association with ENCP. Adjusted Odds Ratio (AOR) with corresponding 95% CI was used to measure the level of the association.

Results

Socio-demographic Characteristics of the Respondent

About 233 health professionals participated in the study making 100% response rate. Less than half, 99(42.5%) of the respondents were in the age ranges of 26-30 years. The mean (\pm SD) ages of the respondents were 27.24 (\pm 3.58) years. Almost half of the respondents 115(49.4%) were married, 95(40.8%) unmarried, 15(6.4%) divorced, and 8(3.4%) were widowed. More than half of them were male 123(52.8%) (Table 1).

Table 1 Socio-Demographic Characteristics of Health Professionals Providing Essential Newborn Care Practice in Rural Gedeo Zone Health Facilities, Southern Ethiopia, 2017 ($N = 233$)

Variables	Frequency (number)	Percentage (%)
Religion		
Orthodox Christians	105	45.2
Protestant	102	43.7
Muslims	26	11.1
Professions		
Midwifery	75	32.2
Health officer	42	18
Nurse	116	49.8
Place of work		
Health center	206	88.4
Hospital	27	11.6
Work experience		
1-5 years	194	83.4
6-10 years	30	12.8
Above 10 years	9	3.8
Monthly income (USD)		
53.74-76.87	66	28.3
76.87-100.12	81	34.8
100.12- 123.38	60	25.7
123.38- 146.64	13	5.6
> 146.64	13	5.6
Qualification		
Diploma	147	63.1
Degree	86	36.9
Institution they attained education		
Private institution	77	33
Government institution	156	67

Essential Newborn Care Practice

Results from observation of the obstetric room at the commencement of labour and delivery, 83.7% were clean. Majority, 222 (95.5%) of health professionals arranged cord cutter and clasper before the start of delivery (Table 2). Almost three fourth, 173(74.3%) of them did not prepare new born identification band; 209(89.7%) of health professionals equipped suction device before the start of delivery. Moreover, 221(94.8%) of health professionals prepared neonatal Ambu

bag and mask before the onset of delivery, but only 75 (32.2%) of them provide vitamin *K* for the newborn, and 177(76%) of them did not place newborn's identification band on the wrist and ankle.

Above half, 138(59.2%) of them practiced hand washing with water and soap and dried by using cloths; most of them, 217(93.1%) put on the sterile glove, and 186(79.8%) of the health professionals wipes the eye and face of the newborn

Table 2 The Level of Essential Newborn Care Practice among Professionals in Rural Gedeo Zone Health Facilities, SNNPR, Southern Ethiopia, 2017 ($N = 233$)

Essential Newborn care practice observed	Frequency	Percentage (%)
Immediately dry the whole body, while assessing the baby's breathing		
Yes	177	76
No	56	24
Evaluate breathing		
Yes	225	96.6
No	8	3.4
Cord care		
Yes	224	96.1
No	9	3.9
Kangaroo mother care (KMC)		
Yes	58	24.9
No	175	75.1
Helps to initiate immediate breast milk feeding within 1 hour		
Yes	189	81.1
No	44	18.9
Administer Eye drop/TTC eye ointment		
Yes	161	69.1
No	72	30.9
Administer Vitamin K injection		
Yes	75	32.2
No	158	67.8
Place the newborn's identification bands on the wrist and ankle		
Yes	56	24
No	177	76
Weigh the newborn		
Yes	190	81.5
No	43	18.5
Vaccination at birth(OPV0 and BCG) given		
Yes	177	76
No	56	24

Note: *OPV*: oral polio virus vaccine, *BCG*: Bacillie Calmete Guarin vaccine

when the head is delivered; 58(24.6%) of them cleaned eyes of the newborn after birth by using a separate swab., and 205(87.6%) of health workers directly dried the whole body with cloth/towel. Fifty-eight (44.9%) of health professionals kept warm by placing the baby skin-to-skin contact or kangaroo mother care (KMC); 211(90.6%) of the workers protected the baby's body and head with a hygienic cloth, and all most all health professionals, 225 (96.5%) checked whether the baby is crying while drying. Most of health professionals 161(69.1%) administered Eye drop/TTC eye ointment and 75(32.2%) administer Vitamin *K* injection. Over all, among the total of 233 respondents, 177(76%) had poor newborn care practice and the rest 56(24%) had good newborn care practice.

Knowledge of Respondents about Newborn Care

Among 233 participants, 47 (20.2) of health workers had good knowledge, 146 (62.7) had fair knowledge and 40 (17.1) of health workers were not knowledgeable (scored below the mean). Regarding the steps of the newborn care, above half, 127 (54.5) did not know all the 10 steps, less half of them, 106 (45.5) knew the 10 steps of newborn care practice.

Health care professionals were requested about newborn care practice steps and 92.7% of health care workers said that they immediately put newborn on mother's abdomen; 91 responded that they dry baby and give eye care. 83.7% of the respondents reported that they assess breathing; 88.8% reported that they practice cord cutting and care; 81.5% of them said they make early initiation of breast feeding. Moreover, 76% of the respondents reported that they practice skin to skin contact; 78.5% provide TTC eye ointment, and 70% provide vitamin *K*. 76% of the health care professionals provide vaccination and 83.7% weigh the newborn.

Most, 215(92.2%) of health professionals know the complications of immediate newborn,

whereas few of them, 18(7.8%) do not know the complications of immediate newborn. Among those who know the newborn complication, 181(77.7%) of them said the complication is hypothermia, whereas 174(74.7%) said asphyxia. However, 155(66.5%) of respondents said infection are mostly occurred complications of immediate newborn.

Training and Supportive Supervision

Almost half, 114(48.9%) of health professionals got in service training about essential newborn care, from which 70(30%) trained once, 37(15%) trained two times and 7(3.0%) trained three times. About 97(41.6%) of them got supportive supervision from their higher officials and nongovernmental organizations (NGOs).

Factors associated with essential newborn care practice

Female respondents were three times more likely to provide good newborn care practice (NBCP) than respondents who are male [adjusted odds ratio (AOR) = 3.08; 95% CI (1.07, 8.81)]. Participants who are in the age group between 36 and 40 years were 81.8 less likely to provide good newborn care as compared to those between 21 and 25 years [AOR = 0.18; 95 CI (0.03, 0.89)]. Health care workers who have got supportive supervision within the last 3 months were 2 times more likely to give good newborn care than those health care workers who haven't got supportive supervision [AOR = 2.17; 95% CI (1.12, 4.19)].

Health care workers who got training related to newborn care were four times more likely to offer good newborn care practice than their counterparts [AOR = 4.25; 95% CI (2.33, 10.09)].

The odds of respondents who have knowledge about the complication of newborn are two times more likely to provide good newborn care practice than the odds of respondents who have no knowledge about the complication of newborn [AOR = 1.98; 95% CI (1.00,3.89)] (Table 3).

Table 3 The factors associated with essential newborn care practice among health professionals of rural Gedeo zone health facilities, Southern Ethiopia, February, 2017 (n=233)

List of Variables and Responses	ENCP		COR with 95% CI	AOR with 95% CI
	Good	Poor		
Age in years				
21-25	13	62	1	1
26-30	16	83	5.56(1.60,19.29)	1.37(0.50,3.73)
31-35	12	20	6.05(1.79,20.39)	0.75(0.20,2.76)
36-40	8	6	1.94(0.52,7.16)	0.18(0.03, 0.89)*
>40	7	6	0.87(0.19,3.99)	0.18(0.03,1.12)
Sex of respondents				
Male	36	87	1	1
Female	20	90	1.86(1.00,3.46)	3.08(1.07,8.81)*
Religion				
Orthodox	24	81	1	1
Protestant	23	79	1.01(0.53,1.9)	2.60(0.85,7.94)
Muslim	9	17	0.56(0.22,1.41)	0.76(.16,3.55)
Monthly Salary in (USD)				
53.74-76.87	12	54	1	1
76.87-100.12	22	59	7.87(1.98,31.25)	0.92(0.26,3.22)
100.12- 123.38	9	51	4.69(1.25,17.61)	0.50(0.11,2.22)
123.38-146.64	6	7	8.11(2.02,32.59)	3.44(0.22,52.95)
> 146.64	7	6	3.93(0.71, 21.59)	3.50(0.16,72.3)
Training status				
Trained	42	72	4.37(2.22,8.59)	4.25(2.33,10.09)*
Not trained	14	105	1	1
Respondent's work experience				
< 5 years	36	158	1	1
5-10 years	13	13	2.19(0.52,9.19)	0.09(0.01, 0.60)
> ten years	7	6	0.38(0.08,1.82)	1.14(0.16,7.89)
Status of supportive supervision				
Yes	32	65	2.297(1.24,4.23)	2.17(1.12,4.19)*
No	24	112	1	1
knowledge on ENCP				
Yes	17	89	2.32(1.22,4.40)	0.43(1.22,3.81)
No	39	88	1	1
Knowledge on Complication				
Yes	40	148	2.04(1.0,4.12)	1.98(1.00,3.89)*
No	16	29	1	1

Note: * significantly associated factors ($P<0.05$)

Discussion

A good newborn care is fundamental for the existence of newborn baby [15]. It includes essential basic newborn care practices (*i.e.* ensuring the air way is clear and remove mucus from the mouth, nose and throat with suction, immediate kangaroo mother care to prevent hypothermia, clean and dry newborn with warm towel, clamp and cut umbilical cord with sterile instrument, providing eye care with tetracycline eye ointment or silver nitrate, giving vitamin *K* and vaccine, weigh newborn and start breast feeding within one hour) [6].

The result of the current study disclosed that only 24% of health care workers were practicing good ENBCP (newborn care that achieves all the elements of newborn care). This finding is in line with other literatures in southern Ethiopia, Chenchu district [16]. However, the result of this study presented that the level of ENCP in Gedeo zone health facilities is lower than a study done in Jimma (51.1%) [11], Addis Ababa (80.7%) [17], Tigray (72.77%) [12], and Khartoum (41.1%) [18]. These differences may be due to study sites in which the current study was conducted in rural health facilities where human resource, medication and materials needed for immediate newborn care practice are limited [19,20].

Respondents providing good ENCP among female professionals were three times higher as compared to male professionals. This might be due to the higher commitment and motherhood feeling of females for newborns and children [21,22].

The odds of giving newborn care decrease by 81.8% among health professionals who are within the age ranges of 36-40 years as compared to 21-25 years. This result is supported by a study done at Indonesia [23]. The possible explanation might be due to the older health professionals are less likely to access updated procedures of essential newborn care practice, and profession-

als may become tiresome for their work as they become older [24].

Participants who had got supportive supervision in the last 3 months were 2.2 times more likely to deliver good ENCP as compared to those who had not got supportive supervision. This confirms the idea that supportive supervision has a greater value for health professionals by motivating and updating of essential new born care procedures and protocols [25].

The odds of providing good ENCP among respondents attended newborn care related training were 4.2 times higher than those who never attended training. This conclusion is supported by another similar study done in Tigray [11]. This might be due to the attitudinal change, gain of knowledge and experience sharing while attending professional trainings [26,27].

Limitation

There might be Hawthorns effect, and since it is a cross sectional study design, it might not display the direct cause and effects association of dependent and independent variables.

Conclusion

The overall essential newborn care practices among health professionals were not good. Age range of 30-40 years, female sex, supportive supervision, attending training, and knowing complication of the newborn were significantly associated factors with essential newborn care practice. This reveals a requirement to strengthen training and supportive supervision platforms to health professionals working in the rural health centers by the Zonal health department and district health offices, and non-governmental organizations.

Assertions

Abbreviations and Acronyms

AOR	Adjusted Odds Ratio
EDHS	Ethiopian Demographic Health Survey
ENAP	Every Newborn Action Plan
ENCP	Essential New Born Care Practice
IRB	Institutional Ethical Review Board
KMC	Kangaroo Mother Care
MEDHS	Mini Ethiopian Demographic Health Survey
NGOs	Non-Governmental Organizations
OR	Odds Ratio
OPV0	Oral Poliovirus Vaccine
QOC	Quality of Care
SPSS	Software Package For Social Science
SNNPR	Southern Nation, Nationalities Peoples Region
USD	United States Dollar

Acknowledgments

Our deepest heart felt gratitude goes to Dilla University for providing the chance and giving ethical clearance. We thank Gedeo zone health Bureau and Gedeo zone health centers for their collaboration. We acknowledged data collectors, supervisors and all study respondents who spent their fruitful time for the accomplishment of this study.

Ethical Concern

Before the actual data collection, ethical clearance was obtained from Dilla University institutional ethical review board (IRB), and support letter was accessed from Gedeo Zone Health office. Each respondent gave written consent after clarification about the scope and purposes of the study. The participants were up-to-date about the right to decline and or to discontinue their involvement at any time they want and no harm will be enforced due to their discontinuation. The filled questionnaires were approved for its completeness and uniformity by the investigators and supervisors. The challenges were solved on time through discussion with concerned bodies. Appropriate explanation/correction was given for a professional found having unfavorable knowledge and or who misses ENCP steps during the data collection.

Consent for Publication - Not applicable

Availability of Data and Materials

The datasets underlying the study are available from the corresponding author up on request.

Competing Interests

We confirm that there are no competing interests on this research work.

Funding

This study has received funding from Dilla University, NORHED Project I.

Author's Detail

¹Department of Reproductive Health, School of Public health, Medical and Health Science College, Dilla University, Dilla, Ethiopia.

²Department of Public Health, School of Public health, Medical and Health Science College, Dilla University, Dilla, Ethiopia.

³Department of Psychiatry, Medical and Health Science College, Dilla University, Dilla, Ethiopia

References

1. Ashraf N. Clinico-Biochemical Profile in Neonates with Birth Asphyxia. *Journal of Islamabad Medical & Dental College*. 2017; **6**(2):64-8.
2. Callaghan-Koru JA, Seifu A, Tholandi M, de Graft-Johnson J, Daniel E, Rawlins B, et al. Newborn care practices at home and in health facilities in 4 regions of Ethiopia. *BMC pediatrics*. 2013; **13**(1):198.
3. Mekonnen Y, Tensou B, Telake DS, Degefe T, Bekele A. Neonatal mortality in Ethiopia: trends and determinants. *BMC public health*. 2013; **13**(1):483.
4. Basavanthappa B. Textbook of midwifery and reproductive health Nursing: Jaypee Brothers Publishers; 2006.
5. Haines A, Sanders D, Lehmann U, Rowe AK, Lawn JE, Jan S, et al. Achieving child survival goals: potential contribution of community health workers. *The Lancet*. 2007; **369**(9579):2121-31.
6. Bezerra FD, Menezes MA, Mendes RB, Santos JM, Leite DC, Kassab SB, Gurgel RQ. Perinatal care in a northeastern Brazilian state: structure, work processes, and evaluation of the components of essential newborn care. *Revista Paulista de Pediatria*. 2019 Apr; **37**(2):140-8.
7. Slattery MM, Morrison JJ. Preterm delivery. *The Lancet*. 2002 Nov 9; **360**(9344):1489-97.
8. Renfrew MJ, McFadden A, Bastos MH, Campbell J, Channon AA, Cheung NF, et al. Midwifery and quality care: findings from a new evidence-informed framework for maternal and newborn care. *The Lancet*. 2014; **384**(9948):1129-45.
9. Waiswa P, Peterson S, Tomson G, Pariyo GW. Poor newborn care practices-a population based survey in eastern Uganda. *BMC pregnancy and childbirth*. 2010; **10**(1):9.

10. Thaver D, Zaidi AK. Burden of neonatal infections in developing countries: a review of evidence from community-based studies. *The Pediatric infectious disease journal*. 2009; **28**(1):S3-S9.
11. Negussie B, Hailu F, Megenta A. Knowledge and Practice of Essential Newborn Care and Associated Factors among Nurses and Midwives Working at Health Centers in Jimma Zone, Ethiopia, 2016. *J Nurs Care*. 2018; **7**(446):2167-1168.1000446.
12. Berhe AK, Tinsae F, Gebreegziabher G. Knowledge and practice of immediate newborn care among health care providers in eastern zone public health facilities, Tigray, Ethiopia, 2016. *BMC pediatrics*. 2017; **17**(1):157.
13. Tsegay T. Knowledge and Practice of Immediate Newborn Care among Midwives in Governmental Health Facilities at Central Zone, Tigray Regional State, North Ethiopia: Addis Ababa University; 2015.
14. Chomba E, McClure EM, Wright LL, Carlo WA, Chakraborty H, Harris H. Effect of WHO newborn care training on neonatal mortality by education. *Ambulatory Pediatrics*. 2008; **8**(5):300-4.
15. Kuhn P, Sizun J, Casper C, GREEN study group from the French Neonatal Society, Allen A, Audoud F, Bouvard C, Brandicourt A, Cayemaex L, Denoual H, Duboz MA. Recommendations on the environment for hospitalised newborn infants from the French neonatal society: rationale, methods and first recommendation on neonatal intensive care unit design. *Acta Paediatrica*. 2018 Nov; **107**(11):1860-6.
16. Mersha A, Assefa N, Teji K, Shibiru S, Darghawth R, Bante A. Essential newborn care practice and its predictors among mother who delivered within the past six months in Chench District, Southern Ethiopia, 2017. *PloS one*. 2018; **13**(12):e0208984.
17. Wondaferash MT. Assessment of knowledge, attitude, and practice about immediate newborn care among health care providers in Addis Ababa Public Health centers. *Ethiopian Journal of Pediatrics and Child Health*. 2011; **7**(7):42-79.
18. NasorTaha FA. Assessment of knowledge, Attitude and Practices of nurse midwives towards immediate care of the newborn in Khartoum state teaching hospitals (2011). *Journal of American Science*. 2013; **9**(9).
19. Dynes M, Buffington ST, Carpenter M, Handley A, Kelley M, Tadesse L, et al. Strengthening maternal and newborn health in rural Ethiopia: Early results from frontline health worker community maternal and newborn health training. *Midwifery*. 2013; **29**(3):251-9.
20. Chopra M, Lawn JE, Sanders D, Barron P, Karim SSA, Bradshaw D, et al. Achieving the health Millennium Development Goals for South Africa: challenges and priorities. *The Lancet*. 2009; **374**(9694):1023-31.
21. Baxter J. Employment characteristics and transitions of mothers in the Longitudinal Study of Australian Children. Australian Government Department of Social Services Occasional Paper. 2013 Dec 18(50).
22. Kachel AF, Premo LS, Hublin JJ. Grandmothering and natural selection. Proceedings of the Royal Society B: *Biological Sciences*. 2011 Feb 7; **278**(1704):384-91.
23. Sethi R, Tholandi M, Amelia D, Pedrana A, Ahmed S. Assessment of knowledge of evidence-based maternal and newborn care practices among midwives and nurses in six provinces in Indonesia. *International Journal of Gynecology & Obstetrics*. 2019; **144**:51-8.
24. Hirani S. Child-rearing practices in Pakistan and associated challenges for health care professionals. Global Unity for Neonatal Nurses(Online). 2008.
25. Otolorin E, Gomez P, Currie S, Thapa K, Dao B. Essential basic and emergency obstetric and newborn care: from education and training to service delivery and quality of care. *International Journal of Gynecology & Obstetrics*. 2015; **130**:S46-S53.
26. Opiyo N, English M. In-service training for health professionals to improve care of the seriously ill newborn or child in low and middle-income countries. Cochrane database of systematic reviews. 2010(4).
27. Senarath U, Fernando DN, Rodrigo I. Effect of training for care providers on practice of essential newborn care in hospitals in Sri Lanka. *Journal of Obstetric, Gynecologic & Neonatal Nursing*. 2007; **36**(6):531-41.