



Backyard Sheep Fattening: Constraints, and Economic Implications in Bule District, Gedeo Zone, Ethiopia

Tsegaye Undesa Galcha¹ and Wondewsen Bekele Wondatir^{2*} 

¹Bule Woreda Agricultural Office, Livestock And Fisher Development, Bule, Ethiopia;

²Department of Animal and Range Sciences, Dilla University; P.O.Box.33, Dilla, Ethiopia;

*Corresponding author; Email: wondewsen19@gmail.com

Received: 10 March 2025

Accepted: 5 May 2025

Published: 30 June 2025

©2025 Dilla University. All Rights Reserved

Article DOI:10.20372/ejed.v07i1.04

Abstract

The research was carried out in the Bule district of the Gedeo Zone to evaluate the practices, challenges, opportunities, and economic feasibility of backyard sheep fattening. Six out of the 18 Kebeles were intentionally selected based on the availability of sheep resources, the experience of farmers, and ease of access. A total of 126 households were randomly selected for the survey. Sheep fattening was widespread in the region, with 94.4% of households employing tethering and free-grazing methods. Traditional fattening techniques were predominant, with an average of 1.97 ± 0.08 sheep being fattened over durations ranging from 2 to 6 months, and 1.59 ± 0.059 cycles per year. Rams were the preferred choice for fattening (72.2%), followed by a combination of rams and ewes (17.5%), and ewes alone (10.3%). The Easter season (27.78%) was the most common time for fattening, and 40.48% of fatteners engaged in this practice primarily for income generation. Separate housing for sheep was common, although often poorly managed; 35.7% of households housed sheep separately within other livestock shelters. Key challenges included feed shortages, diseases, rising veterinary drug prices, decreasing grazing land, low sheep productivity, and issues with predators and theft. Conversely, opportunities included favorable agro-ecological conditions, abundant fresh grass, experienced farmers, a large sheep population, increasing market demand, rising sheep prices, higher literacy rates, and larger household sizes, all of which could enhance fattening practices. Despite these opportunities, sheep fattening remained unprofitable under current management systems. To improve profitability, the study recommends targeted training, ongoing support, collaboration among stakeholders, improved forage development, and consistent monitoring and evaluation.

Keywords/Phrases: Backyard, Bule, Easter, Fattening, Profitability, Ram

1 Introduction

Ethiopia boasts one of the most substantial livestock populations in Africa, comprising around 70.3 million cattle, 42.9 million sheep, 52.5 million goats, 8.2 million camels, 2 million horses, 9 million donkeys, 0.38 million mules, and 49 million chickens (CSA, 2021). For many rural communities, livelihoods are deeply intertwined with livestock production, complementing traditional agricultural practices (Estefanos *et al.*, 2014). Sheep, in particular, play a vital role in Ethiopian livestock, being exten-

sively reared across the country (Tadesse *et al.*, 2015; Abera and Elias, 2019). As a result, sheep fattening has emerged as a crucial economic activity that enhances food security and supports vulnerable groups, including women and disadvantaged members of society (Teketel *et al.*, 2021).

Despite the significance of sheep in the local economy, productivity faces numerous challenges (Tariku & Etefa, 2022). While previous studies have attempted to address these issues, most have focused on specific feeding systems within controlled envi-

ronments (Skunmun *et al.*, 2012; Wude, 2017; Gebrekidan, 2018; Sime, 2019; Kokeb *et al.*, 2021; Alemu, 2023). This focus highlights a critical gap in research regarding farmers using diverse feeding systems that utilize local feed resources.

Bule District in the Gedeo Zone is recognized as a potential hub for sheep production, benefiting from a favorable climate and a rich tradition of backyard sheep fattening. The district features a robust livestock market and hosts a diverse sheep population, including various breeds from the Oromia and Sidama regions. However, modern sheep fattening practices remain largely absent, and economic growth among sheep fatteners has been sluggish. There is a notable lack of cooperative associations, investors, or initiatives aimed at developing sheep fattening into a viable business. Many farmers treat sheep fattening as a traditional activity rather than a reliable income source, leading to inconsistent practices and a reliance on basic grazing without additional management strategies.

Furthermore, a comprehensive investigation of backyard sheep fattening, including constraints, opportunities, and profitability in this area, has not yet been conducted. Current evidence suggests that in-

adequate research and limited technological skills in fattening practices prevent households from capitalizing on potential economic benefits throughout the sheep value chain (Wamatu *et al.*, 2022). Therefore, this study aims to assess backyard sheep fattening practices, identify existing limitations, explore potential opportunities, and evaluate the economic profitability of sheep fattening in Bule District.

2 Materials and methods

2.1 Description of the study area

The research area is located in the Gedeo Zone of the South Ethiopia Regional State, roughly 386 km away from Addis Ababa. It covers an area of 27,300 hectares and has an elevation range between 2,001 and 3,000 meters above sea level. The average annual precipitation in this region is approximately 1,600 mm, with temperatures varying from 12.6°C to 20°C. The main agricultural system practiced here is a mixed crop-livestock production system. Major crops include barley, beans, peas, maize, and wheat, along with perennial plants such as enset and coffee. The area borders the Oromiya region to the south, east, and west, and the Sidama Zone to the north (Bule Woreda BOPE, 2022, unpublished).




Figure 1. Study area map

2.2 Sampling method and sample size determination

The research utilized a multistage sampling technique. Initially, six kebeles (Bule 01, Bule 03, Sika, Kochor, Suqo, and Igna Okolu) out of a total of 18 were purposefully chosen based on the availability of sheep resources, the experience of farmers in sheep production, and their accessibility. These kebeles account for 33.3% of the total kebeles in the district.

To initiate the selection process, a comprehensive list of all household heads engaged in sheep production was compiled to establish the sampling frame. Representative households were subsequently chosen at random through either simple or systematic random sampling methods, ensuring that key criteria pertinent to sheep fattening practices were adequately represented.

The study focused on sheep-fattening farmers within the selected kebeles, identified from a total population of 328 households as per district livestock records. A sample size of 126 households was calculated using Yamane's formula (1973), applying a standard error of 0.07 and a confidence level of 95%. This led to the selection of 21 households from each of the six kebeles, which were randomly chosen for interviews and data collection.

2.3 Method of data collection

Data were gathered through the use of a semi-structured questionnaire, field observations, key informant interviews, and focus group discussions.

2.3.1. Questionnaire survey

Data was gathered through pre-tested, semi-structured questionnaires that were administered by trained enumerators to a total of 126 randomly selected households. The survey gathered information on fattening practices (cycle, season, sources, and feed conservation) as well as household demographics (age, sex, family size, education, marital status, landholding, and income sources), fattening constraints, and economic profitability.

2.3.2. Field observation

During the administration of the questionnaire, the husbandry methods for fattened sheep—including

management, feeding, and housing conditions—were observed. Additionally, preferred morphological characteristics of the fattening sheep, such as coat color, body length and height, horn shape, tail type, health status, and age determined by dentition, were important components considered during field observations (Kostera, 2007; Ciesielska & Jemielniak, 2018).

2.3.3. Key informant interview

Key informant interviews will be conducted with individuals who possess in-depth knowledge and experience related to backyard sheep fattening. This may include agricultural extension workers, livestock experts, community leaders, cooperative heads, and experienced sheep farmers. The aim is to gather qualitative insights on local practices, challenges, and support systems associated with sheep fattening.

2.3.4. Focus group discussions

Six focus groups were conducted across the selected kebeles, comprising diverse participants, including women and elders. Discussions explored feeding systems, fattening challenges, animal husbandry practices, and potential opportunities for sheep fattening in the region. The insights gathered provided valuable qualitative data to complement the quantitative findings of the study.

2.4 Partial budget analysis

Sheep fattening costs and returns have been estimated using budgeting procedures. Specifically, the budgeting method will make use of gross margin analysis, which is exactly how model's net income was calculated (Rahman & Lawal, 2003).

$$\text{Gross margin} = \text{Gross income} - \text{Total variable cost} \quad (1)$$

$$\text{Net farm income} = \text{Gross margin} - \text{Total fixed cost} \quad (2)$$

2.5 Statistical data analysis

The survey data were examined utilizing SPSS version 20. Descriptive statistics, including mean, per-

centage, and standard deviation, were employed to encapsulate the results. According to a formula of index employed by Musa *et al.*, 2006), indexes were obtained to provide a ranking of the specifically favorable fattening season, mostly preferred type of ram and ewe, fattening sheep selection criteria. Therefore, the index was calculated as,

Index = Sum of $[(r_n * C_1 + r_{n-1} * C_2 + \dots + r_1 * C_n)]$ raised for an individual season, preference, and criteria divided by the sum of $[(r_n * C_1 + r_{n-1} * C_2 + \dots + r_1)]$ for the overall season, preference and criteria.

Where,

r_n = Value for the least ranked level (for example if the least rank is 5th rank, then $r_{n-5}, r_{n-1} = 4$, and... $r_1 = 1$); C_n = Counts of the least ranked level (in the above example, the count of the 5th rank = C_n , and the counts of the 1st rank = C_1).

3 Results and Discussions

3.1 Demographic characteristics of households

In the study area, respondents had an average family size exceeding the national rural average of 4.9 members per household (Ayele & Gebretsadik, 2024). This increase can be attributed to cultural influences that view family members as valuable resources for labor in agricultural activities and social status, as well as the prevalence of polygamous marriages, often pursued to acquire additional farmland (see Table 1).

The average age of sheep fatteners in the study area was 41.38 years, ranging from 19 to 83 years old. This finding aligns with Zemene & Tolemariam's (2016) report of 41.9 years for households in the Jimma Zone. However, it is lower than the average

ages reported by Teklay (2008) and Endale (2015), who found averages of 48.1 and 49.14 years, respectively. The age range is comparable to Yeshitila's (2008) study in Alaba, which reported an average of 42.82 years. This age distribution suggests that an active and potentially interested population is engaged in sheep fattening and broader agricultural activities within the study area.

Households in the study area had an average total landholding of 0.76 hectares, significantly less than the 1.93 hectares reported in the South Gondar Zone (Tilahun, 2021). Fortunately, literacy rates are promising, with 76.3% of sheep fatteners having at least some formal education, which could facilitate the adoption of improved farming techniques.

Group discussions revealed a local tradition that dictates livestock ownership in the area belongs to the husband, even if the wife has her own fattening sheep. This finding is consistent with Gebru *et al.* (2017), who reported that, similar to other African nations, household resource leaders are primarily men.

3.2 Occupation of the households

In the study area, pure agriculture dominated household occupations, accounting for 87.3% of fatteners. This was followed by agriculture combined with small trade (6.3%), agriculture with labor work (2.4%), and agriculture combined with civil service (4%). The dominance of pure farming, which typically involves mixed crop cultivation and livestock rearing, can likely be attributed to favorable agro-ecological conditions that support both agricultural and livestock activities. This observation aligns with findings from West Hararghe, Oromia, where mixed crop-livestock farming serves as the primary livelihood and income source (Abdi *et al.*, 2013).

Table 1. Fattener socio-economic data in study area

Variables		Kebeles						Total N=126	p-value		
FS	Mean ± SE	5.62 ± .71 ^b	5.10 ± .46 ^c	5.43 ± .50 ^b	5.14±.46 ^c	7.14±.43 ^a	7.43±.62 ^a	5.98±.23	0.004		
	Mi	1	3	2	2	4	3	1			
	Ma	14	10	12	10	10	13	14			
Age	Mean ± SE	36.24 ± 2.81 ^c	37.33 ± 1.6 ^c	45.90 ± 3.26 ^a	40.05 ± 1.64 ^b	42.00±1.8 ^b	46.76±2.6 ^a	41.38±1.01	0.007		
	Mi	19	24	26	25	28	29	19			
	Ma	72	46	83	56	56	81	83			
TLH(ha)	Mean ± SE	0.70 ± .071 ^b	.69 ± .058 ^c	0.67 ± .053 ^c	0.851 ± .087 ^a	0.76±.098 ^b	0.90±.06 ^a	0.76±.035	0.094		
	Mi	0.23	0.25	0.25	0.25	0.25	0.35	.023			
	Ma	1.65	1	1.03	1.70	1.25	1.67	1.70			
MS	Married%	14.3	16.7	14.3	15.9	15.9	15.9	92.9	0.610		
	Single%	0.8	0.0	1.6	0.0	0.0	0.0	2.4			
	Widowed%	1.8	0.0	0.8	0.8	0.8	0.8	4.8			
Sex of Respondents (%)											
Education level of HHs		Male (n=108)		Female (n=18)		Total	χ^2	p-value			
		Illiterate		25.9		77.8	33.3	18.051	0.001		
		1-4		34.3		5.6	30.2				
		5-8		29.6		11.1	27.0				
		9-10		5.6		0.0	4.8				
		Diploma		2.8		5.6	3.2				
		Degree		1.9		0.0	1.6				
Occupation of fattener/sources of income				Frequency (N=126)			%				
				Agriculture (crop and livestock production)			110				
							87.3				
				Agriculture and petty-trade			8				
							6.3				
				Agriculture and labour work			3				
							2.4				
				Agriculture and civil servant			5				
							4.0				

SE=standard error Ma = maximum; Mi = minimum; FS = family size; TLH = total land hold; MS = marital status; HH = households; N = number of respondents.
 The above table adopted from Tsegaye and Wondewesen (2024) which was published on Online Journal of Animal and Feed Research

3.3 Number, Duration and fattening cycle

On average, fatteners in the study area completed 1.50 ± 0.050 sheep fattening cycles per year, ranging from 1 to 3 rounds, with no significant differences across kebeles. The findings of the current study differ from those of Shewangzaw *et al.* (2018) in the Amhara region, where the average number of fattened sheep per fattening period was higher at 2.86. However, these findings are consistent with those of Assefa & Ayza (2020) in the Hadiya Zone of Southern Ethiopia, where farmers typically practice fattening 2 to 3 times a year.

In contrast, Nurlign (2020) reported predominantly 2 cycles, while Getachew *et al.* (2017) noted only one cycle in the Fogera District. Samuel (2016) found similar results of 1.625 ± 0.05 rounds per year in the Amhara region. The study suggests that improving feed and nutritional status, as well as management practices, may be key to increasing both the number of fattened sheep per cycle and the total number of fattening cycles per year. The variation in the number of fattening cycles across study kebeles may be attributed to differences in resources, agro-ecological conditions, and seasonal variations in fattening (see Table 2).

Table 2. Number, duration and cycle of fattening sheep in study area

Variables		Kebeles						Overall N=126	p-value
		Bule 01	Bule03	sika	suquo	kochore	1gna okolu		
NFS	Mean \pm SE	1.6 \pm .14 ^c	1.4 \pm .13 ^c	2.05 \pm .18 ^a	2.2 \pm .22 ^a	2.1 \pm .24 ^a	2.1 \pm .22 ^a	1.9 \pm .08 ^b	0.02
	Mi	1	1	1	1	1	1	1	
	Ma	3	3	4	4	5	4	5	
FD	Means	4.1 \pm .96 ^a	4 \pm .13 ^a	3.9 \pm .19 ^b	3.8 \pm .19 ^b	3.6 \pm .16 ^c	3.6 \pm .16	3.8 \pm .07 ^c	0.2
	Mi	2	3	2	2	2	2	2	
	Ma	6	5	5	5	5	5	6	
FC	Mean \pm SE	1.8 \pm .159 ^a	1.3 \pm .126 ^c	1.4 \pm .11 ^c	1.7 \pm .15 ^a	1.5 \pm .14 ^b	1.5 \pm .14 ^b	1.5 \pm .05	0.09
	Mi	1	1	1	1	1	1	1	
	Ma	3	3	2	3	3	3	3	

NFS = Number of fattening sheep FD = Fattening duration; FC = Fattening cycle; N = total sampled respondents; Mi = Minimum; Ma Maximum; SE = standard error

3.4 Fatteners ranking criteria for selection of sheep fattening

In the study area, sheep fatteners employed specific selection criteria, prioritizing conformation (height, length, and appearance) through visual assessment (see Table 3). This criterion was ranked as the most important factor (Index = 0.22), followed closely by physical characteristics such as color, horn shape, and tail type (Index = 0.21). Age and health status were equally important, ranking third (Index = 0.17), while adaptability and price were considered less crucial.

These findings align with previous research (ESG-PIP, 2012; Assefa & Ayza, 2020; Diriba *et al.*, 2021; Getahun, 2022), which emphasizes factors like body condition, frame, age, and health. Fatteners primarily relied on visual cues, such as body frame, hair appearance, and the presence of diarrhea, to assess sheep suitability. While most respondents (65.1%) considered age important, they relied on visual estimation rather than dentition; however, a significant portion (34.9%) deemed age less critical for fattening purposes.

Table 3. Fattening Sheep Selection Criteria

No	Selection Criteria	Frequency of rank given fatteners						Total	Index	Rank
		1 st	2 nd	3 rd	4 th	5 th	6 th			
1	Body conformation (height, length and appearance)	55	9	34	16	13	3	130	0.22	1
2	Physical characteristics	11	64	23	18	5	2	123	0.21	2
3	Age	46	11	3	1	26	42	129	0.17	3
4	Health	9	23	32	32	27	6	129	0.17	3
5	Price	4	2	13	11	33	61	124	0.09	6
6	Adaptability	1	17	21	48	22	12	121	0.14	5
Mostly preferred physical characteristics										
1	Color	22	39	21	20	-	-	102	0.27	2
2	Horn	15	24	37	33	-	-	109	0.24	3
3	Tail	5	16	33	39	-	-	93	0.17	4
4	Body length and height	58	21	9	8	-	-	96	0.32	1
Respondent %								Total	χ^2	p-value
Consideration of age of fattening sheep		Bule 01	Bule03	Sika	Suqo	Kochore	1gna okolu		22.071	0.001
Yes (%)		6.3	10.3	14.3	15.9	10.3	7.9	65.1		
No (%)		10.3	6.3	2.4	0.8	6.3	8.7	34.9		

Index = [(6 × number of responses for 1st rank + 5 × number of responses for 2nd rank + 4 × number of responses for 3rd rank + 3 × number of responses for 4th rank + 2 × number of responses for 5th rank + 1 × number of responses for 6th rank)] divided by (6 × total responses for 1st rank + 5 × total responses for 2nd rank + 4 × total responses for 3rd rank + 3 × total responses for 4th rank + 2 × total responses for 5th rank + 1 × number of responses for 6th rank).

3.5 Preferences of fatteners for sheep fattening

The study revealed that the majority (72.2%) of respondents preferred fattening rams due to market demand, feed efficiency, potential profit, and superior weight gain, consistent with findings in the Genji district (Diriba *et al.*, 2021). Mature rams were ranked highest (index value 0.37) because of their rapid growth, high market demand, and favorable pricing, followed by young rams (0.31) and old rams (0.30). Fatteners prioritized mature rams for their quicker attainment of market weight (47.9%), strong market demand (39.6%), and higher selling prices (12.5%), as indicated in Table 4.

In the study area, barren ewes were preferred for fattening due to their marketability, associated with good body condition and non-reproductive use (Index = 0.40). Old ewes were also favored for their low purchase price and suitability for culling (50%) and high demand from local butchers (37.5%). Gimmer ewes were preferred due to market demand and usage by local butchers (Mamo, 2020; Getahun, 2022). This aligns with findings in other regions, where unproductive female sheep or aged animals are commonly fattened due to market factors (Diriba *et al.*, 2021; Samuel, 2016) (see Table 5).

Table 4. Condition of castration practices, preferred fattening sheep type

Variable		Frequency (n)	Percent (%)
Castration practices	Yes	0	0.0
	No	126	100.0
Type of sheep for fattening	Ram	91	72.2
	Ewe	13	10.3
	Both	22	17.5
Total		126	100.0


Table 5. Community preference of fatten sheep in the study area

Mostly preferred type of ram	1 st ranked F	2 nd ranked F	3 rd ranked F	Total	Index	Rank
old ram	23	50	40	113	0.30	3
matured ram	48	47	18	113	0.37	1
young ram	42	16	55	113	0.31	2
Mostly preferred type of Ewe						
Old ewe	8	14	13	35	0.29	2
Barren/infertile ewe	21	13	1	35	0.40	1
Gimmer ewe	6	8	21	35	0.25	3

Index = $[(3 \times \text{number of responses for 1st rank} + 2 \times \text{number of responses for 2nd rank} + 1 \times \text{number of responses for 3rd rank}) \text{ divided by } (3 \times \text{total responses for 1st rank} + 2 \times \text{total responses for 2nd rank} + 1 \times \text{total responses for 3rd rank})]$; F=frequency.

3.6 Sheep fattening occasions

Sheep fattening in the study area is a seasonal practice, with 27.78% of farmers targeting the Easter season (see Fig. 1). This finding aligns with research by Wude (2017) in the Debre-Berhan area and Assefa & Ayza (2020), who also noted peak fattening periods around Easter, Christmas, and the Ethiopian New Year.

**Figure 2.** Seasons of sheep fattening in study area

3.7 Reasons for fattening sheep

The primary motivation for sheep fattening in the study area is income generation, accounting for 40.48%. Other reasons include saving, acquiring additional sheep, and meeting socio-cultural needs. This observation is consistent with the studies con-

ducted by Fikru & Gebeyew (2015) in the Degehabur Zone and Samuel (2016) in the Amhara Region, both of which identified income generation as the principal aim of sheep fattening (refer to Fig. 2).




Figure 3. Main purpose of involving in sheep fattening activities

3.8 Fattening sheep, feeding system and supplementary feed types

The feeding systems and feed types typically utilized for fattening sheep in the study region are detailed in Table 6. The main feeding systems observed are tethering combined with stall feeding (68.3%), semi-grazing alongside stall feeding (20.6%), and free grazing utilizing a cut-and-carry method (11.1%). This result aligns with Getachew *et al.* (2017),

who reported that stall feeding (56.1%), free grazing (18.1%), mixed systems (11.1%), and tethering (7.4%) were common among respondents in the upper land of Fogera District. The current findings also agree with Esatu *et al.* (2019), who noted that tethering, cutting, and carrying were practiced feeding systems for sheep in Arba Minch Zuria District.

Table 6. Provided feed and feeding/fattening system in study area

Description	Sheep fattening/Feeding system (%)			Total	χ^2	p-value
	FGCC	TSFS	SGSFS			
Feed for sheep during stall-feeding	B.L & HL	2.4	20.6	6.3	29.4	
	GFG	4.8	23.8	6.3	34.9	7.779
	CF	4.0	10.3	2.4	16.7	0.4
	Enset	0.0	10.3	3.2	13.5	
	CRB	0.0	3.2	2.4	5.6	
Total		11.1	68.3	20.6	100.0	

B.L & HL = Bamboo leaf & home leftover; GFG = Green fresh grass; CF = Cultivated forage; CRB = crop residues with bole; FGCC = Free grazing with cut and carry system; TSFS = Tethering with stall feeding system; SGSFS = semi-grazing with stall feeding system.

Backyard sheep fatteners in the Bule district commonly tether sheep on nearby pastures, supplementing their diet with fresh grass, bamboo leaves, and enset. Tethering with stall feeding is practiced, with fresh grass (23.8%), bamboo leaves (20.6%), and cultivated forage (10.3%) being the primary feeds. This observation aligns with findings from Salo *et al.* (2017) in Anelemo Woreda, where farmers also tether animals and provide pasture grown between enset.

Tethering is a common sheep fattening practice in the study area, promoting weight gain through grazing, reducing aggression, and preventing breeding (Getachew *et al.*, 2017). This method optimizes feed utilization, reduces labor, and minimizes unnecessary movement, ultimately leading to faster attainment of market weight. Households utilized three main feeding systems: semi-grazing with stall feeding, tethering with stall feeding, and free grazing with cut-and-carry. Semi-grazing with stall feeding involves allowing sheep to graze freely before confining them to a fenced pasture for stall feeding (Esatu *et al.*, 2019). Free grazing with cut-and-carry was also common, where sheep grazed freely for 11 hours a day and were provided with additional supplements (Abebe *et al.*, 2021).

The study found no significant difference ($p > 0.05$) in the types of feed provided to fattening sheep across different feeding systems in the study area. Fresh green grass (34.9%) was the primary feed, supplemented with bamboo leaves and home leftovers, with variations based on kebele and availability. The similarity in feed types across systems and the unknown daily amounts given may explain this finding, with basal feed resources used as supplements during grazing. This aligns with Machen (2019), who also reported variations in supplementary feed types across locations due to factors such as land availability (see Table 7).

3.9 Utilization and type of available supplementary feed

As indicated in Figure 3, supplementary feeds for sheep fattening in the Bule district primarily included salt, kitchen leftovers (such as cereal screenings and coffee residues), and enset (18.3%). Combinations of salt, bole, and home leftovers (15.1%) as well as

enset with salt (11.1%) were also common. Coffee residues served as a readily available daily supplement due to household production. This study aligns with Nurlign (2020), who reported the use of food leftovers and cereal screenings in Ethiopian sheep fattening.

Sheep fatteners in the study area faced significant constraints due to feed shortages. According to household respondents, the primary consequences included weight loss in fattening sheep (47.6%), a reduction in the annual fattening cycle (42.9%), and increased mortality rates among sheep due to harsh conditions (9.5%) (see Table 10). To mitigate these challenges, farmers proposed several measures: purchasing forage or renting grazing land (42.9%), utilizing available crop residues (9.5%), and excluding pasture areas from livestock access (47.6%).

These findings align with results reported by Guyo (2016), who observed that in Bonke Woreda of Gamo Gofa Zone, 62.5% of households in highland areas used farm residues, while 37.5% of mid-altitude producers relied on purchased feed to combat feed shortages. However, the proportions of crop residue users and feed purchasers in the current study were notably lower.

Farmers adopted various strategies before and during feed shortages, including purchasing materials like enset and its residues and renting grazing pastures specifically for fattening animals. This proactive approach indicates that some farmers possess the knowledge and skills needed to anticipate feed shortages and implement preventive measures. Furthermore, extending improved forage cultivation could become viable if supported by training and attention from stakeholders.

Within the study area, some farmers utilized crop residues treated with salt or bole soaking, and 9.5% believed that using crop residues could help mitigate feed shortages. Abshir (2018) similarly emphasized the importance of crop residues in addressing acute feed resource shortages. According to Tolera *et al.* (2012), crop residues account for approximately 50% of the total feed supplied in Ethiopia. Nonetheless, the use of these techniques by fatteners in the current area remains minimal due to factors like low palatability, labor intensity, and time constraints. Respon-

dents indicated that chopping crop residues requires additional labor and time, further complicating their use.

Moreover, Alemu *et al.* (1991) highlighted that the feeding value of crop residues is limited by poor voluntary intake, low digestibility, and deficiencies in nitrogen, energy, minerals, and vitamins. Fatteners noted that population growth and intensified crop cultivation have resulted in insufficient grazing land, compelling them to restrict livestock access to pastures until grass is established for grazing during designated seasons.

To improve the intake and nutritional status of avail-

able feeds, fatteners could enhance utilization efficiency by employing techniques such as chopping, salting, and mixing bole (mineral soil) into feed materials. Among these practices, salting was noted as particularly common; it often followed the chopping of feed into manageable sizes to enhance intake. Sampled households affirmed that employing these techniques significantly alleviated some consequences arising from feed shortages. Supporting evidence from the central highlands of Ethiopia suggests that feed processing methods like chopping, wetting, grinding, boiling, and roasting could significantly contribute to resolving feed-related issues (Ahmed *et al.*, 2010).




Figure 4. Percentage HHs supplementary feeds used

3.10 Economic analysis of sheep fattening

Budgeting analysis of sheep fattening in the Bule district revealed that while farmers achieved a gross margin of 517.3 ETB per sheep after 2 to 6 months of fattening, the practice was ultimately unprofitable. Costs were categorized into fixed expenses (such as equipment depreciation and land rent) and variable costs (including feed and healthcare), with sheep procurement accounting for 61% of total expenses. Feed represented the largest cost component, averaging 694.7 ETB, while variable feeding costs constituted

86.4% of total feed expenditure.

The average cost of purchasing sheep was 2,554.07 ETB, with a selling price of 4,122.23 ETB. The resulting benefit-cost ratio of 0.98 indicates that for every 1 ETB invested, the return was only 0.98 ETB, confirming economic unprofitability under current management practices, as a profitable business typically has a benefit-cost ratio greater than one (Sarma *et al.*, 2014).

This loss is exacerbated by a lack of awareness

among farmers regarding the tracking of variable costs, leading them to primarily focus on the purchase-sale price differential while neglecting other significant input costs. To improve profitability, it is crucial for fatteners to meticulously record all costs associated with sheep fattening, enabling a more comprehensive understanding of the true economic benefits and potential for optimization.

4 Conclusion and Recommendation

4.1 Conclusion

The study identifies challenges in backyard sheep fattening, including inadequate housing, poor feed management, and disease outbreaks. Opportunities exist to improve productivity but require coordinated efforts.

4.2 Recommendations

- Enhance farmer's capacity through training related to proper management practices, including record-keeping and disease control.
- Improve feed access- Develop strategies to improve feed quality and availability, including supplementary feeding and nutrient-rich basal feeds.
- Support from authorities- Provide government support through subsidies, grants, or low-interest loans, and expand training in modern techniques and disease control.
- Encourage best practices- Foster collaboration among stakeholders to implement best practices and raise awareness among farmers about efficient management.
- Promote long-term profitability- Encourage better practices through education and demonstration of long-term profitability.

Acknowledgments

I would like to extend my heartfelt appreciation to the Bule Woreda Agricultural Office, the Livestock and Fishery Office, and the department staff, as well as the dedicated farmers and Agricultural Development Agents for their invaluable support throughout this research project. Additionally, I am deeply grateful to the laboratory staff of the Hawassa and Dilla

University Nutrition Laboratories for their expertise and commitment. Your hard work and professionalism have significantly contributed to the quality and rigor of this research.

Data availability

Data will be made available on request.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this manuscript.

References

Abdi, E., Kamal, K., Yassin, E. and Muleta, D. (2013). Cattle production in west hararge and opportunity and constraints assessments in Darulabou, Gemicheis and Chiro districts Oromiya Regional State Ethiopia. *International Journal of Livestock Production Research*, 1(1): 1- 15.

Abebe, Y., Tefere, M., Shiferaw Dagnew, Y. G., Sellaissie, E. E., Smaling, E. M., & Teshome, A. (2021). Challenges of Shifting from Free Grazing to Adoption of Cut-and-Carry Feeding System in North-Western Ethiopian Highlands. ESAP P, 255.

Abera Anja and Elias Hesebo (2019). Sheep Fattening Practices and Constraints in East Badewacho Woreda, Hadiya Zone Southern Ethiopia. *Global Veterinaria*, 21 (3): 109-115.

Alemu Gashe Desta (2023). Nutritional Content Analysis of Crop Residues in Three Agroecologies in East Gojjam Zone. *Research Article. The Scientific World Journal*. Volume 2023, Article ID 1974081, 8 pages. <https://doi.org/10.1155/2023/1974081>.

Assefa, A., & Ayza, A. (2020). Assessment of sheep fattening and marketing systems in Duna Woreda, Hadiya zone, Southern Ethiopia. *J Dairy Vet Anim Res*, 9(4), 130-135.

Ayele, T., & Gebretsadik, T. (2024). Gender analysis in selected agricultural practices in Gedeo and Halaba zones in Southern Ethiopia. *International*

Journal of Agricultural Research, Innovation and Technology (IJARIT), 14(1), 1-9.

Ciesielska, M., & Jemielniak, D. (Eds.). (2018). *Qualitative methodologies in organization studies* (p. 33). Cham, Switzerland: Palgrave Macmillan.

CSA (2021). Crop yield for major grain crops, private smallholder farms, main season. Addis Ababa, Ethiopia.

Diriba, T., Zemene, W., Solomon, D., and Monenus, E., (2021). Assessment of sheep fattening practices under smallholder farmers in Genji district of West Wollega, Western Oromia, Ethiopia. *International Journal of Veterinary Sciences and Animal Husbandry*; 6(4): 18-25.

Disassa H, Teklu T, Adem H. (2013). A study on ovine pneumonic pasteurellosis: isolation and identification of Pasteurellae and their antibiogram susceptibility pattern in Haramaya District, Eastern Hararghe, Ethiopia. *BMC Vet Res*. 9:1–8. <https://doi.org/10.1186/1746-6148-9-239>.

Endale, Y., (2015). Assessment of Feed Resources and Determination of Mineral Status of Livestock Feed in Meta Robi District, West Shewa Zone, Oromia Regional State, Ethiopia. Msc. Thesis Submitted to the School of Graduate Studies of Ambo University, Ethiopia.

Esatu, B., Yisehak, K., Chencha, C., (2019). Feed resources availability, feeding system and feed balance of sheep in Arba Minch Zuria District of Gamo Zone, Southern Ethiopia. *Research Article. OMO Int. J. Sci. Vol:2 Issue :2 :59-75; (ISSN: 2520 – 4882)*.

ESGPIP (Ethiopia sheep and goat productivity improvement program) (2012). Short Term Intensive Fattening of Sheep and Goats for Rapid Improvement in Weight and Condition and also Producer Incomes, Technical Bulletin No.11.p2-6.

Estefanos T., Tesfaye A, Feyisa H, Gashaye W, Tatek W, Tesfaye K, Osho T. (2014). Traditional cattle production in the highlands of Hararge: Case study for East and West Zones of the high lands of Hararghe, Eastern Ethiopia. *Basic Res.J. Agricult. Sci. Rev.*, 3(12): 122-130.

Fikru, S., & Gebeyew, K. (2015). Sheep and goat production systems in Degehabur Zone, Eastern Ethiopia: Challenge and opportunities. *J Adv Dairy Res*, 3(134), 2.

Gebrekidan, T. (2018).sheep and goat husbandry practices in eastern tigray, and effects of substitution of mulberry (*Morus alba*) leaf meal for concentrate mix on growth and carcass traits of Tigray highland lambs in barley straw based feeding. PhD Dissertation. Addis Ababa University, Addis Ababa, Ethiopia.

Gebru, H., Mohammed, A., Dechassa, N., & Belew, D. (2017). Assessment of production practices of smallholder potato (*Solanum tuberosum L.*) farmers in Wolaita zone, southern Ethiopia. *Agriculture & Food Security*, 6(1), 1-11.

Getachew, M., Firew, T., and Yeshambel, M. (2017). Sheep fattening practices in Fogera District, Amhara National Regional State, Ethiopia. *Journal of Biology, Agri. and Health care*; 7(15):33-39.16.

Getahun, B. (2022). Sheep fattening and marketing systems in Angollelana Tera district, North Shewa Zone, Ethiopia. Msc thesis. Debre Berhan University, Debre Berhan, Ethiopia.

Kokey, M., Mekonnen, Y., & Tefera, M. (2021). Feed intake, digestibility and growth performance of Dorper-Menz cross bred sheep fed local brewery by-product (atella) and concentrate mixture at different levels of combination. *African Journal of Biological Sciences*; 3(4):33–41.

Kostera, M. (2007). *Organizational Ethnography: Methods and Inspirations*. London: Sage.

Machen, R. (2019). Supplementation of grazing sheep. Texas A&M Agrilife Ext., TX, USA. Available at:(accessed 2 July 2020).

Mamo, B., (2020). Assessment of Sheep Fattening Practices around Chalia District West Shoa Zone. *J. Ethol & Animal Sci*, 3(1): 000116.

Musa, L. M. A., K. J. Peters and M. K. A. Ahmed, (2006). On farm characterization of Butana and Kenana cattle breed production systems in Sudan. *Livest. Res. for Rural Develop.* 18 (12).

Nurlign, M. (2020). Sheep Fattening, Marketing Systems and Constraints of Ethiopia: A Review. *World Appl. Sci. J.*, 38 (5): 416-421.

Rahman, S. A., & Lawal, A. B. (2003). Economic analysis of maize-based cropping systems in Giwa Local Government Area of Kaduna State, Nigeria. *An International Journal of Agricultural Sciences, Science, Environment and Technology*, 3(2), 139-148.

Salo, S., Tadesse, G., & Haylemeskel, D. (2017). Survey on constraints of improved forage adoption in Anelemo Woreda, Hadiya Zone, Ethiopia. *Agri Cultural Research and Technology: Open Access Journal*, 12(2).

Samuel, M. (2016). Characterization of sheep fattening at small holder level in different Agro-Ecological Zones of Ethiopia–Amhara Region. M.Sc. Thesis. Bahir Dar University, Bahir Dar, Ethiopia.

Sarma P. K., S. K. Raha and H. Jørgensen (2014). An economic analysis of beef cattle fattening in selected areas of Pabna and Sirajgonj Districts. *J. Bangladesh Agril. Univ.* 12(1): 127–134.

Shewangzaw M. A., Teshager, A. A., Endeshaw, A. G., Birhan Atinaw, M., & Tesfa Sendeku, A. (2018). Small ruminant fattening practices in Amhara region, Ethiopia. *Agriculture & Food Security*, 7: 64.

Sime, T. E. (2019). Assessment of production status and evaluation of biomass yield, chemical composition and supplementary value of mulberry (*Morus alba*) on growth performance and carcass characteristics of yearling arsi-bale sheep (Doctoral dissertation).

Skunmun, P., Boonyanuwat, P.K., Koonavoot, R.S. and Suwanasopee, T. (2012). Improving Smallholder and Industrial Livestock Production for Enhancing Food Security, Environmental and human welfare. Proceedings of the 15th AAAP Animal Science Congress held at Thammasat University, Pathum Thani, 26-30, November 2012, AVAT. Pp. 267.

Tadesse, E., Negesse, T., Abebe, G. (2015). Sheep production and marketing system in southern Ethiopia: the case of Awassa zuria district. *Trop. Anim. Health Prot.* 47(7):1417-1425.

Tariku, B., & Etefa, M. (2022). Study on Improved Sheep Fattening Production, Value Addition and Marketing in Bekiyo Bekit Kebele, Adiyo District, Kaffa Zone, Southwest Ethiopia. *Journal of Applied Agricultural Economics and Policy Analysis*, 5(1), 9-18.

Teketel, M., Temesgen, D., Hamza, D., & Fesseha, H. (2021). Determinants of smallholder farmers' participation in improved sheep production: The case of Doyogena District, Kembata Tembaro Zone, Southern Ethiopia. *Advances in Agriculture*, 2021, 1-17.

Teklay, A. (2008). Assessment of the Feeding Systems and Feed Resources of Dairy Cattle in Lemu-Bilbilo Wereda Dairy Products-Processing Cooperatives, Arsi Zone of Oromia Regional State, Ethiopia. Addis Ababa University, Addis, Ethiopia.

Tilahun, D. (2021). Assessment of available livestock feed resources in South Gondar Zone, Amhara National Regional State, Ethiopia. *American Journal of Agriculture and Forestry*, 9, 269-275.

Wamatu, J., Ekule, M., & Mebratie, A. (2022). Assessment of knowledge, attitude and practice of smallholder farmers on market-oriented sheep fattening in North Shewa, Ethiopia.

Wude T. (2017). Sheep production systems in urban and peri-urban areas of debre-berhan and dessie, ethiopia, and performance of lambs fed different combinations of wheat bran and lentil screening. PhD Dissertation. Addis Ababa University, Addis Ababa, Ethiopia.

Yamane, Taro. (1973). *Statistics and Introductory analysis*, 3rd Ed., New York: Harper and Row.

Yeshitila, A. (2008). Assessment of Livestock Feed Resources Utilization In Alaba Woreda, Southern Ethiopia. Msc Thesis. Haramya University, Haramya, Ethiopia.

Zemene, W., Tilahun, S., & Tolemariam, T. (2016). Assessment of the prevailing cattle fattening practices in Jimma Zone, South-Western Ethiopia. *Global Veterinaria*, 17(2), 105-113.